
Product Name: KD-Validated EGFR Recombinant Rabbit Monoclonal Antibody**Catalog #: KVA01002**

For research use only.

Summary

Description	KO&KD-Validated antibody
Host	Rabbit
Application	WB,FCM,ICC,IHC-P
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Rabbit IgG
Clonality	Rabbit mAb
Form	Liquid
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1,000-1:5,000; FC 1:200-1:2,000; ICC 1:100-1:1,000; IHC-P 1:100-1:200
Molecular Weight	Calculated MW: 134.3kDa

Antigen Information

Gene Name	EGFR EGFR; Epidermal Growth Factor Receptor; ERBB1; ERBB; Receptor Tyrosine-Protein Kinase ErbB-1; Erb-B2 Receptor Tyrosine Kinase 1; Proto-Oncogene C-ErbB-1; EC 2.7.10.1; HER1; Epidermal Growth Factor Receptor (Avian Erythroblastic Leukemia Viral (V-Erb-B)
Alternative Names	Oncogene Homolog); Erythroblastic Leukemia Viral (V-Erb-B) Oncogene Homolog (Avian); Avian Erythroblastic Leukemia Viral (V-Erb-B) Oncogene Homolog; Epidermal Growth Factor Receptor Tyrosine Kinase Domain; Cell Proliferation-Inducing Protein 61; Cell Growth Inhibiting Protein 40; EGFR VIII; EC 2.7.10; NISBD2; PIG61; MENA
Gene ID	1956.0
SwissProt ID	P00533
Immunogen	A synthesized peptide derived from human EGFR




Background


The protein encoded by this gene is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor, thus inducing receptor dimerization and tyrosine autophosphorylation leading to cell proliferation. Mutations in this gene are associated with lung cancer. EGFR is a component of the cytokine storm which contributes to a severe form of Coronavirus Disease 2019 (COVID-19) resulting from infection with severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2). [provided by RefSeq, Jul 2020]

Research Area


Signal Transduction,Cell Biology,Microbiology,Cancer,Neuroscience

Image Data







Immunocytochemical staining of HepG2 cells using EGFR antibody (KVA01002, 1:1,000), Top panel: wild-type (WT); Bottom panel: EGFR shRNA knockdown (KD). Nuclei were stained blue with DAPI; EGFR was stained magenta with Alexa Fluor® 647. Scale bar, 20 μm. Permeabilization: Triton.



Immunocytochemical staining of HepG2 cells with EGFR antibody (KVA01002, 1:1,000). Nuclei were stained blue with DAPI; EGFR was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser intensity and smart gain: Medium. Scale bar: 20 μm.



Validation of EGFR knockdown using flow cytometry. Wild-type (WT, Blue) and knockdown (KD, Green) HepG2 cells were stained with EGFR antibody (KVA01002, 1:2,000) and analyzed using BD flow cytometer.



Flow cytometric analysis of EGFR expression in HepG2 cells using EGFR antibody (KVA01002, 1:2,000). Green, isotype control; red, EGFR.