

Product Name: KD-Validated TAR DNA Binding Protein Recombinant Rabbit Monoclonal Antibody

Catalog #: KVA01474

For research use only.

Summary

Description	KO&KD-Validated antibody
Host	Rabbit
Application	WB,FCM,ICC
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Rabbit IgG
Clonality	Rabbit mAb
Form	Liquid
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1,000-1:5,000; FC 1:200-1:2,000; ICC 1:100-1:1,000
Molecular Weight	Calculated MW: 44.7kDa

Antigen Information

Gene Name	TARDBP
Alternative Names	TARDBP; TAR DNA Binding Protein; TDP-43; ALS10; TAR DNA-Binding Protein 43; TAR DNA-Binding Protein-43; TDP43
Gene ID	23435.0
SwissProt ID	Q13148
Immunogen	A synthesized peptide derived from human TDP43

Background


HIV-1, the causative agent of acquired immunodeficiency syndrome (AIDS), contains an RNA genome that produces a chromosomally integrated DNA during the replicative cycle. Activation of HIV-1 gene expression by the transactivator Tat is

dependent on an RNA regulatory element (TAR) located downstream of the transcription initiation site. The protein encoded by this gene is a transcriptional repressor that binds to chromosomally integrated TAR DNA and represses HIV-1 transcription. In addition, this protein regulates alternate splicing of the CFTR gene. A similar pseudogene is present on chromosome 20. [provided by RefSeq, Jul 2008]


Research Area

Microbiology, Neuroscience, Epigenetics and Nuclear Signaling


Image Data




Western blotting analysis using TAR DNA binding protein antibody (KVA01474). TAR DNA binding protein expression in wild type (WT) and TAR DNA binding protein (TARDBP) shRNA knockdown (KD) HeLa cells with 30 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with TAR DNA binding protein antibody (KVA01474, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Western blotting analysis using TAR DNA binding protein antibody (KVA01474). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with TAR DNA binding protein antibody (KVA01474, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Immunocytochemical staining of C2C12 cells with TAR DNA binding protein antibody (KVA01474, 1:1,000). Nuclei were stained blue with DAPI; TAR DNA binding protein was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar: 20 µm.



Flow cytometric analysis of TAR DNA binding protein expression in C2C12 cells using TAR DNA binding protein antibody (KVA01474, 1:2,000). Green, isotype control; red, TAR DNA binding protein.