
Product Name: KD-Validated Phospho-c-Myc (S62) Recombinant Rabbit Monoclonal Antibody**Catalog #: KVA01538**

For research use only.

Summary

Description	KO&KD-Validated antibody
Host	Rabbit
Application	WB,FCM,ICC
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Rabbit IgG
Clonality	Rabbit mAb
Form	Liquid
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1,000-1:5,000; FC 1:200-1:2,000; ICC 1:100-1:1,000
Molecular Weight	Calculated MW: 48.8kDa

Antigen Information

Gene Name	MYC MYC Proto-Oncogene, BHLH Transcription Factor; BHLHe39; C-Myc; MYCC; V-Myc Avian Myelocytomatosis Viral Oncogene Homolog; Class E Basic Helix-Loop-Helix Protein 39; Myc
Alternative Names	Proto-Oncogene Protein; Transcription Factor P64; Proto-Oncogene C-Myc; Myc-Related Translation/Localization Regulatory Factor; Avian Myelocytomatosis Viral Oncogene Homolog; V-Myc Myelocytomatosis Viral Oncogene Homolog; BHLHE39; MRTL
Gene ID	4609.0
SwissProt ID	P01106
Immunogen	A synthesized peptide derived from human Phospho-c-Myc (S62)


Background

This gene is a proto-oncogene and encodes a nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. The encoded protein forms a heterodimer with the related transcription factor MAX. This complex binds to the E box DNA consensus sequence and regulates the transcription of specific target genes. Amplification of this gene is frequently observed in numerous human cancers. Translocations involving this gene are associated with Burkitt lymphoma and multiple myeloma in human patients. There is evidence to show that translation initiates both from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site, resulting in the production of two isoforms with distinct N-termini. [provided by RefSeq, Aug 2017]


Research Area

Epigenetics and Nuclear Signaling, Stem Cells, Cancer, Signal Transduction


Image Data




Western blotting analysis using phospho-c-Myc (S62) antibody (KVA01538). Phospho-c-Myc (S62) expression in wild type (WT) and MYC knockdown (KD) HSHC cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with phospho-c-Myc (S62) antibody (KVA01538, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Western blotting analysis using phospho-c-Myc (S62) antibody (KVA01538). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with phospho-c-Myc (S62) antibody (KVA01538, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Immunocytochemical staining of HeLa cells with Phospho-c-Myc (S62) antibody (KVA01538, 1:1,000). Nuclei were stained blue with DAPI; Phospho-c-Myc (S62) was stained magenta with Alexa Fluor® 647. Images were taken using Leica stellaris 5. Protein abundance based on laser Intensity and smart gain: Medium. Scale bar, 20 µm.



Flow cytometric analysis of Phospho-c-Myc (S62) expression in HeLa cells using Phospho-c-Myc (S62) antibody (KVA01538, 1:2,000). Green, isotype control; red, Phospho-c-Myc (S62).