

Product Name: KD-Validated Phospho-EGFR (Y1068) Recombinant Rabbit Monoclonal Antibody

Catalog #: KVA01737

For research use only.

Summary

Description	KO&KD-Validated antibody
Host	Rabbit
Application	WB,FCM,ICC,IHC-P
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Rabbit IgG
Clonality	Rabbit mAb
Form	Liquid
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Supplied in PBS (pH 7.4) containing 50% glycerol, and 0.02% sodium azide.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:1,000-1:5,000; FC 1:200-1:2,000; ICC 1:100-1:1,000; IHC-P 1:100-1:200
Molecular Weight	Calculated MW: 134.3kDa

Antigen Information

Gene Name	EGFR EGFR; Epidermal Growth Factor Receptor; ERBB1; ERBB; Receptor Tyrosine-Protein Kinase ErbB-1; Erb-B2 Receptor Tyrosine Kinase 1; Proto-Oncogene C-ErbB-1; EC 2.7.10.1; HER1; Epidermal Growth Factor Receptor (Avian Erythroblastic Leukemia Viral (V-Erb-B) Oncogene Homolog); Erythroblastic Leukemia Viral (V-Erb-B) Oncogene Homolog (Avian); Avian Erythroblastic Leukemia Viral (V-Erb-B) Oncogene Homolog; Epidermal Growth Factor Receptor Tyrosine Kinase Domain; Cell Proliferation-Inducing Protein 61; Cell Growth Inhibiting Protein 40; EGFR VIII; EC 2.7.10; NISBD2; PIG61; MENA
Gene ID	1956.0
SwissProt ID	P00533

Immunogen

A synthesized peptide derived from human Phospho-EGFR (Y1068)


Background

The protein encoded by this gene is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor, thus inducing receptor dimerization and tyrosine autophosphorylation leading to cell proliferation. Mutations in this gene are associated with lung cancer. EGFR is a component of the cytokine storm which contributes to a severe form of Coronavirus Disease 2019 (COVID-19) resulting from infection with severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2). [provided by RefSeq, Jul 2020]


Research Area

Signal Transduction, Cell Biology, Microbiology, Cancer, Neuroscience


Image Data




Western blotting analysis using phospho-EGFR (Y1068) antibody (KVA01737). Phospho-EGFR (Y1068) expression in wild type (WT) and EGFR shRNA knockdown (KD) HepG2 cells with 20 µg of total cell lysates. Hsp90 α serves as a loading control. The blot was incubated with phospho-EGFR (Y1068) antibody (KVA01737, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.




Western blotting analysis using phospho-EGFR (Y1068) antibody (KVA01737). Total cell lysates (30 µg) from various cell lines were loaded and separated by SDS-PAGE. The blot was incubated with phospho-EGFR (Y1068) antibody (KVA01737, 1:5,000) and HRP-conjugated goat anti-rabbit secondary antibody (APS0635, 1:10,000) respectively.



Immunohistochemistry was performed on paraffin-embedded human lung adenocarcinoma using phospho-EGFR (Y1068) antibody (KVA01737, 1:200). Antigen retrieval was done in sodium citrate buffer (pH 6.0). DAB was used for detection, with hematoxylin counterstaining. Images were acquired using a Nikon Ci-L Plus microscope (40× objective). Scale bar: 25 µm.



Immunocytochemical staining of HepG2 cells using Phospho-EGFR (Y1068) antibody (KVA01737, 1:1,000), Top panel: wild-type (WT); Bottom panel: Phospho-EGFR (Y1068) shRNA knockdown (KD). Nuclei were stained blue with DAPI; Phospho-EGFR (Y1068) was stained magenta with Alexa Fluor® 647. Scale bar, 20 µm. Permeabilization: Triton.



Validation of EGFR knockdown using flow cytometry. Wild-type(WT, Blue) and knockdown(KD, Green) HepG2 cells were stained with Phospho-EGFR (Y1068) antibody (KVA01737, 1:2,000) and analyzed using BD flow cytometer.