
Product Name: CD4 (7H9) Mouse Monoclonal Antibody**Catalog #: AMM00731**

For research use only.

Summary

Description	Mouse monoclonal Antibody
Host	Mouse
Application	IHC
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG1
Clonality	Monoclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% sodium azide, pH 7.3.
Purification	Affinity Purification

Application

Dilution Ratio	IHC 1:50-1:100
Molecular Weight	-

Antigen Information

Gene Name	CD4
Alternative Names	CD4; T-cell surface glycoprotein CD4; T-cell surface antigen T4/Leu-3; CD antigen CD4
Gene ID	920
SwissProt ID	P01730
Immunogen	Synthetic Peptide of CD4

Background


Cluster of Differentiation 4 (CD4) is a glycoprotein composed of an amino-terminal extracellular domain (four domains: D1-D4 with Ig-like structures), a transmembrane part and a short cytoplasmic tail. CD4 is expressed on the surface of T helper cells,

regulatory T cells, monocytes, macrophages and dendritic cells, and plays an important role in the development and activation of T cells. On T cells, CD4 is the co-receptor for the T cell receptor (TCR), and these two distinct structures recognize the Antigen–Major Histocompatibility Complex (MHC).


Research Area

Immunology


Image Data




Immunohistochemistry analysis of paraffin-embedded Human stomach tissue using CD4 (7H9) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Human tonsils using CD4 (7H9) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Immunohistochemistry analysis of paraffin-embedded mouse brain tissue using CD4 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.



Immunofluorescence analysis of CD4 (7H9) in mouse colon tissue using CD4 (7H9) antibody (11A1) (red), and DAPI (blue).



Immunofluorescence analysis of CD4 (7H9) in rat lung using CD4 antibody (11A1) (red), and DAPI (blue).



Immunohistochemistry analysis of paraffin-embedded Human Amygdala using CD4 (7H9) antibody. High-pressure and temperature Tris-EDTA pH 8.0 was used for antigen retrieval.