

Product Name: Cystatin C (6F12) Mouse Monoclonal Antibody**Catalog #: AMM03309**

For research use only.

Summary

Description	Mouse monoclonal Antibody
Host	Mouse
Application	ELISA
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG1
Clonality	Monoclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% sodium azide, pH 7.3.
Purification	Affinity Purification

Application

Dilution Ratio	ELISA 1:5000-1:20000
Molecular Weight	-

Antigen Information

Gene Name	CST3
Alternative Names	CST3; Cystatin-C; Cystatin-3; Gamma-trace; Neuroendocrine basic polypeptide; Post-gamma-globulin
Gene ID	1471
SwissProt ID	P01034
Immunogen	Recombinant protein of human Cystatin C

Background


Cystatin C is a 14 kDa member of the Cystatin superfamily of cysteine protease inhibitors. Most cell types secrete Cystatin C.

Cystatin C inhibits cathepsins, and thereby may function as a tumor suppressor by inhibiting cathepsin mediated tumor cell invasion. In addition, this tumor suppressor function can also be attributed to Cystatin C's ability to antagonize TGF- β 1 signaling.

Research Area

Tags & Cell Markers

Image Data



Standard Curve for Human Cystatin C: Capture Antibody mouse mAb [6F12C7D8] to Human Cystatin C at 4ug/ml and Detector Antibody mouse mAb [7F6A5F3] to Human Cystatin C at 0.5ug/ml.