

**Product Name:** Phospho-Histone H2A.X (Ser139) (2A9) Mouse Monoclonal Antibody  
**Catalog #:** AMM03679

For research use only.

## Summary

<b>Description</b>	Mouse monoclonal Antibody
<b>Host</b>	Mouse
<b>Application</b>	WB,ICC/IF
<b>Reactivity</b>	Human,Mouse
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Phosphorylated
<b>Isotype</b>	IgG2a
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% sodium azide, pH 7.3.
<b>Purification</b>	Affinity Purification

## Application

<b>Dilution Ratio</b>	WB 1:500-1:1000,ICC/IF 1:50-1:200
<b>Molecular Weight</b>	Calculated MW: 15 kDa; Observed MW: 15 kDa

## Antigen Information

<b>Gene Name</b>	H2AX
<b>Alternative Names</b>	H2A.X; H2AFX; H2a/x; HIST5-2AX; Histone H2A.X; gamma H2A.X
<b>Gene ID</b>	3014
<b>SwissProt ID</b>	P16104
<b>Immunogen</b>	A synthetic Phosphorylated peptide corresponding to residues target protein

## Background


Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a

central role in transcription regulation, DNA repair, DNA replication and chromosomal stability.


## Research Area

Epigenetics and Nuclear Signaling


## Image Data




Western blot analysis of Phosphorylation of H2A.X at Serine 139 in 3T3 or Hydroxyureatreated 3T3 lysates using Phospho-Histone H2A.X (Ser139) antibody.



Immunofluorescence analysis of Phospho-Histone H2A.X (Ser139) (2A9) in 3T3 or Hydroxyureatreated 3T3 using Phospho-Histone H2A.X (Ser139) (2A9) antibody



Immunofluorescence analysis of Phospho-Histone H2A.X (Ser139) (2A9) in A549(upper, untreated or Hydroxyureatreated) and HeLa(lower, untreated or Hydroxyureatreated) using Phospho-Histone H2A.X (Ser139) antibody.



Western blot analysis of Phospho-Histone H2A.X (Ser139) (2A9) in untreated or Hydroxyureatreated HeLa and A549 lysates using Phospho-Histone H2A.X (Ser139) (2A9) antibody(upper) or betaActin antibody(2000688F10) (lower).

