

# **Product Name: Cyclin B1(1A5)Mouse Monoclonal Antibody**

Catalog #: AMM09585

For research use only.

### **Summary**

**Description** Mouse monoclonal Antibody

**Host** Mouse

**Application** IHC,ICC/IF

**Reactivity** Human,Rat,Mouse

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

**Clonality** Monoclonal

Form Liquid Concentration 1mg/ml

**Storage** Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

## **Application**

**Dilution Ratio** IHC 1:100-1:200,ICC/IF 1:50-1:200

Molecular Weight 55kDa

# **Antigen Information**

Gene Name CCNB1
Alternative Names CCNB1
Gene ID 891.0
SwissProt ID P14635

**Immunogen** Synthetic Peptide of Cyclin B1 at AA range of 60-140

# **Background**

The protein encoded by this gene is a regulatory protein involved in mitosis. The gene product complexes with p34(cdc2) to form the maturation-promoting factor (MPF). Two alternative transcripts have been found, a constitutively expressed transcript

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




and a cell cycle-regulated transcript, that is expressed predominantly during G2/M phase. The different transcripts result from the use of alternate transcription initiation sites. [provided by RefSeq, Jul 2008],developmental stage:Accumulates steadily during G2 and is abruptly destroyed at mitosis.,function:Essential for the control of the cell cycle at the G2/M (mitosis) transition.,PTM:Ubiquitinated by the SCF(NIPA) complex during interphase, leading to its destruction. Not ubiquitinated during G2/M phases.,similarity:Belongs to the cyclin family.,similarity:Belongs to the cyclin family. Cyclin AB subfamily.,subunit:Interacts with the CDC2 protein kinase to form a serine/threonine kinase holoenzyme complex also known as maturation promoting factor (MPF). The cyclin subunit imparts substrate specificity to the complex. Binds HEI10. Interacts with catalytically active RALBP1 and CDC2 during mitosis to form an endocytotic complex during interphase.,

#### **Research Area**

 $Cell\_Cycle\_G1S; Cell\_Cycle\_G2M\_DNA; Oocyte\ meiosis; p53; Progesterone-mediated\ oocyte\ maturation;$ 

#### **Image Data**



Immunohistochemical analysis of paraffin-embedded Human lung Tissue using Cyclin B1 Mouse mAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Human Tonsil Tissue using Cyclin B1 Mouse mAb diluted at 1:200.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838