

Product Name: RB Mouse Monoclonal Antibody**Catalog #: AMM80520**

For research use only.

Summary

| | |
|----------------------|---|
| Description | Mouse monoclonal Antibody |
| Host | Mouse |
| Application | IHC,ELISA |
| Reactivity | Human |
| Conjugation | Unconjugated |
| Modification | Unmodified |
| Isotype | Mouse IgG1 |
| Clonality | Monoclonal |
| Form | Liquid |
| Concentration | 1mg/ml |
| Storage | Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles. |
| Shipping | Ice bags |
| Buffer | PBS containing 0.03% sodium azide. |
| Purification | Affinity Purification |

Application

| | |
|-------------------------|---------------------------------------|
| Dilution Ratio | IHC 1:200-1:1000,ELISA 1:5000-1:20000 |
| Molecular Weight | / |

Antigen Information

| | |
|--------------------------|---|
| Gene Name | RB |
| Alternative Names | RB; pRb; OSRC; pp110; p105-Rb; RB1 |
| Gene ID | 5925.0 |
| SwissProt ID | P06400 |
| Immunogen | Purified recombinant fragment of human RB expressed in E. Coli. |


Background

The Rb protein regulates differentiation, apoptosis, and cell cycle control by coordinating the cell cycle at G1-S with transcriptional machinery. During G1, cyclin D-dependent kinase-mediated phosphorylation of Rb at Ser-795 marks the conversion of Rb from a transcriptionally repressive, hypophosphorylated state to an inactive, phosphorylated state, which may

be sustained through mitosis by differential phosphorylation of up to 16 putative serine or threonine residues. Pediatric cancer retinoblastoma and the formation of other human tumors can be attributed to mutations in the retinoblastoma tumor suppressor gene(Rb).

Research Area

Image Data



Immunohistochemical staining of paraffin-embedded human normal esophagus (A) and stomach (B) tissue, showing nucleus localization using Rb mouse mAb with DAB staining.