
Product Name: MAPK10 Mouse Monoclonal Antibody**Catalog #: AMM80648**

For research use only.

Summary

Description	Mouse monoclonal Antibody
Host	Mouse
Application	WB,ICC,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Mouse IgG1
Clonality	Monoclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	PBS containing 0.03% sodium azide.
Purification	Affinity Purification

Application

Dilution Ratio	WB 1:500-1:2000,ICC 1:200-1:1000,ELISA 1:5000-1:20000
Molecular Weight	53kDa

Antigen Information

Gene Name	MAPK10
Alternative Names	JNK3; JNK3A; PRKM10; p54bSAPK
Gene ID	5602.0
SwissProt ID	P53779
Immunogen	Purified recombinant fragment of human MAPK10 (aa28-233) expressed in E. Coli.

Background


MAPK10: mitogen-activated protein kinase 10, also known as JNK3, JNK3A, PRKM10, p54bSAPK. Entrez Protein NP_002744. It is a member of the MAP kinase family. MAP kinases act as an integration point for multiple biochemical signals, and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation and development. This

protein is a neuronal-specific form of c-Jun N-terminal kinases (JNKs). Through its phosphorylation and nuclear localization, this kinase plays regulatory roles in the signaling pathways during neuronal apoptosis. Beta-arrestin 2, a receptor-regulated MAP kinase scaffold protein, is found to interact with, and stimulate the phosphorylation of this kinase by MAP kinase kinase 4 (MKK4). Cyclin-dependent kinase 5 can phosphorylate, and inhibit the activity of this kinase, which may be important in preventing neuronal apoptosis. Four alternatively spliced transcript variants encoding distinct isoforms have been reported.


Research Area

TGF-beta signaling pathway, MAPK signaling pathway

Image Data



Western blot analysis using MAPK10 mouse mAb against NIH/3T3 (1) and SKN-SH (2) cell lysate.



Immunofluorescence staining of methanol-fixed A431 (left) and HeLa (right) cells showing cytoplasmic and membrane localization.