

Product Name: CTNNB1 Mouse Monoclonal Antibody**Catalog #: AMM80788**

For research use only.

Summary

Description	Mouse monoclonal Antibody
Host	Mouse
Application	IHC, ICC, ELISA, FC
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Mouse IgG1
Clonality	Monoclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	PBS containing 0.03% sodium azide.
Purification	Affinity Purification

Application

Dilution Ratio	IHC 1:200-1:1000, ICC 1:200-1:1000, ELISA 1:5000-1:20000, FC 1:200-1:400
Molecular Weight	85kDa

Antigen Information

Gene Name	CTNNB1
Alternative Names	CTNNB; FLJ25606; FLJ37923; DKFZp686D02253; CTNNB1; Beta-catenin
Gene ID	1499.0
SwissProt ID	P35222
Immunogen	Purified recombinant fragment of human CTNNB1 expressed in E. Coli.

Background


The protein encoded by this gene is part of a complex of proteins that constitute adherens junctions (AJs). AJs are necessary for the creation and maintenance of epithelial cell layers by regulating cell growth and adhesion between cells. The encoded protein also anchors the actin cytoskeleton and may be responsible for transmitting the contact inhibition signal that causes

cells to stop dividing once the epithelial sheet is complete. Finally, this protein binds to the product of the APC gene, which is mutated in adenomatous polyposis of the colon. The distinct peripheral cytosolic proteins, alpha, beta and gamma catenin (102, 94 and 86 kDa) are found in many tissues and bind to the conserved cytoplasmic tail domain of the cell adhesion cadherins. Catenins link E cadherin to other integral membrane or cytoplasmic proteins and are modulated by Wnt1 proto oncogene. The central core region of beta catenin is involved in mediation of cadherin catenin complex interaction with EGFR. Beta-Catenin-mediated signalling is involved at several stages of vertebrate neural development.


Research Area

Wnt signaling pathway, Hippo signaling pathway


Image Data



Immunofluorescence analysis of A549 (left) and SK-BR-3 (right) cells using CTNNB1 mouse mAb (green). Red: Actin filaments have been labeled with DY-554 phalloidin. Blue: DRAQ5 fluorescent DNA dye.



Flow cytometric analysis of A549 cells using CTNNB1 mouse mAb (green) and negative control (purple).



Immunohistochemical analysis of paraffin-embedded human Placenta tissues using CTNNB1 mouse mAb