

---

**Product Name: TNNI2 Mouse Monoclonal Antibody****Catalog #: AMM81045**

For research use only.

**Summary**

|                      |   |
|----------------------|---|
| <b>Description</b>   | Mouse monoclonal Antibody   |
| <b>Host</b>          | Mouse   |
| <b>Application</b>   | IHC,ELISA,FC  |
| <b>Reactivity</b>    | Human   |
| <b>Conjugation</b>   | Unconjugated  |
| <b>Modification</b>  | Unmodified  |
| <b>Isotype</b>       | Mouse IgG1  |
| <b>Clonality</b>     | Monoclonal  |
| <b>Form</b>          | Liquid  |
| <b>Concentration</b> | 1mg/ml  |
| <b>Storage</b>       | Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles. |
| <b>Shipping</b>      | Ice bags  |
| <b>Buffer</b>        | Purified antibody in PBS with 0.05% sodium azide.                           |
| <b>Purification</b>  | Affinity Purification   |

**Application**

|                         |  |
|-------------------------|--|
| <b>Dilution Ratio</b>   | IHC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400 |
| <b>Molecular Weight</b> | 21kDa  |

**Antigen Information**

|                          |  |
|--------------------------|--|
| <b>Gene Name</b>         | TNNI2  |
| <b>Alternative Names</b> | DA2B; FSSV; fsTnl; AMCD2B  |
| <b>Gene ID</b>           | 7136.0   |
| <b>SwissProt ID</b>      | P48788   |
| <b>Immunogen</b>         | Purified recombinant fragment of human TNNI2 expressed in E. Coli. |


**Background**

This gene encodes a fast-twitch skeletal muscle protein, a member of the troponin I gene family, and a component of the troponin complex including troponin T, troponin C and troponin I subunits. The troponin complex, along with tropomyosin, is responsible for the calcium-dependent regulation of striated muscle contraction. Mouse studies show that this component is


also present in vascular smooth muscle and may play a role in regulation of smooth muscle function. In addition to muscle tissues, this protein is found in corneal epithelium, cartilage where it is an inhibitor of angiogenesis to inhibit tumor growth and metastasis, and mammary gland where it functions as a co-activator of estrogen receptor-related receptor alpha. This protein also suppresses tumor growth in human ovarian carcinoma. Mutations in this gene cause myopathy and distal arthrogyrosis type 2B. Alternatively spliced transcript variants have been found for this gene.

## Research Area


## Image Data




Red: Control Antigen (100ng); Purple: Antigen (10ng); Green: Antigen (50ng); Blue: Antigen (100ng);




Flow cytometric analysis of NIH/3T3 cells using TNNI2 mouse mAb (blue) and negative control (red).



Immunohistochemical analysis of paraffin-embedded human rabbit cardiac muscle tissues using TNNI2 mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded human cardiac muscle tissues using TNNI2 mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded human striated muscle tissues using TNNI2 mouse mAb with DAB staining.