

Product Name: PRKACA Mouse Monoclonal Antibody**Catalog #: AMM81371**

For research use only.

Summary

Description	Mouse monoclonal Antibody
Host	Mouse
Application	ELISA
Reactivity	Human
Conjugation	Unconjugated
Modification	Unmodified
Isotype	Mouse IgG1
Clonality	Monoclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Purified antibody in PBS with 0.05% sodium azide.
Purification	Affinity Purification

Application

Dilution Ratio	ELISA 1:5000-1:20000
Molecular Weight	40.6kDa

Antigen Information

Gene Name	PRKACA
Alternative Names	PKACA
Gene ID	5566.0
SwissProt ID	P17612
Immunogen	Purified recombinant fragment of human PRKACA (AA: 1-120) expressed in E. Coli.

Background


cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its effects by activating the cAMP-dependent protein kinase, which transduces the signal through phosphorylation of different target proteins. The inactive kinase holoenzyme is a tetramer composed of two regulatory and two catalytic subunits. cAMP causes the dissociation of the

inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. The protein encoded by this gene is a member of the Ser/Thr protein kinase family and is a catalytic subunit of cAMP-dependent protein kinase. Alternatively spliced transcript variants encoding distinct isoforms have been observed.

Research Area

MAPK signaling pathway

Image Data



Black line: Control Antigen (100 ng); Purple line: Antigen(10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng);