# **Product Name: PRDM5 Mouse Monoclonal Antibody**

Catalog #: AMM81582



## **Summary**

Production Name PRDM5 Mouse Monoclonal Antibody

**Description** Mouse Monoclonal Antibody

**Host** Mouse

Application WB,FC,ELISA

**Reactivity** Human

# **Performance**

ConjugationUnconjugatedModificationUnmodifiedIsotypeMouse IgG1ClonalityMonoclonalFormLiquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw

cycles.

**Buffer** Purified antibody in PBS with 0.05% sodium azide

**Purification** Affinity Purification

### **Immunogen**

Storage

Gene Name PRDM5

Alternative Names BCS2; PFM2

Gene ID 11107.0

Q9NQX1.Purified recombinant fragment of human PRDM5 (AA: 1-100) expressed in E.

Coli.

# **Application**

**SwissProt ID** 

**Dilution Ratio** WB:1:500-1:2000,FC:1:200-1:400,ELISA:1:10000

Molecular Weight 73kDa

# **Background**


# Product Name: PRDM5 Mouse Monoclonal Antibody Catalog #: AMM81582




The protein encoded by this gene is a transcription factor of the PR-domain protein family. It contains a PR-domain and multiple zinc finger motifs. Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
<br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription factors of the PR-domain family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription family are known to be involved in cell differentiation and tumorigenesis. <br/>
| Transcription family are known to be involved in cell differentiation and tumorigenesi

#### **Research Area**


### **Image Data**



Black line: Control Antigen (100 ng); Purple line: Antigen(10ng); Blue line: Antigen (50 ng); Red line: Antigen (100 ng);



Western blot analysis using PRDM5 mouse mAb against HL-60 (1) cell lysate.



Flow cytometric analysis of HeLa cells using PRDM5 mouse mAb (green) and negative control (red).

# Product Name: PRDM5 Mouse Monoclonal Antibody

Catalog #: AMM81582



## Note

For research use only.