

# **Product Name: CD274 Mouse Monoclonal Antibody**

Catalog #: AMM82137

For research use only.

### **Summary**

**Description** Mouse monoclonal Antibody

**Host** Mouse

**Application** IHC,ELISA,FC

**Reactivity** Human

ConjugationUnconjugatedModificationUnmodifiedIsotypeMouse IgG2bClonalityMonoclonalFormLiquid

Concentration 1mg/ml

**Storage** Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

**Buffer** Purified antibody in PBS with 0.05% sodium azide

**Purification** Affinity Purification

# **Application**

**Dilution Ratio** IHC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400

Molecular Weight 33.3kDa

# **Antigen Information**

Gene Name CD274

Alternative Names B7-H; B7H1; PDL1; PD-L1; PDCD1L1; PDCD1LG1

 Gene ID
 29126.0

 SwissProt ID
 Q9NZQ7

**Immunogen** Purified recombinant fragment of human CD274 (AA: 24-153) expressed in E. Coli.

# **Background**


This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine




production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants.

#### Research Area


#### **Image Data**



Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)




Flow cytometric analysis of HL-60 cells using CD274 mouse mAb (green) and negative control (red).



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissues using CD274 mouse mAb with DAB staining.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Immunohistochemical analysis of paraffin-embedded human rectum cancer tissues using CD274 mouse mAb with DAB staining.