

---

**Product Name: ATP6V0A4 Mouse Monoclonal Antibody****Catalog #: AMM82886**

For research use only.

**Summary**

<b>Description</b>	Mouse monoclonal Antibody
<b>Host</b>	Mouse
<b>Application</b>	IHC,ELISA,FC
<b>Reactivity</b>	Human
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	Mouse IgG1
<b>Clonality</b>	Monoclonal
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Purified antibody in PBS with 0.05% sodium azide
<b>Purification</b>	Affinity Purification

**Application**

<b>Dilution Ratio</b>	IHC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400
<b>Molecular Weight</b>	96.3kDa

**Antigen Information**

<b>Gene Name</b>	ATP6V0A4
<b>Alternative Names</b>	A4; STV1; VPH1; VPP2; DRTA3; RTA1C; RTADR; ATP6N2; RDRTA2; ATP6N1B
<b>Gene ID</b>	50617.0
<b>SwissProt ID</b>	Q9HBG4
<b>Immunogen</b>	Purified recombinant fragment of human ATP6V0A4 (AA: 228-390) expressed in E. Coli.


**Background**

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of intracellular compartments of eukaryotic cells. V-ATPase dependent acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-


ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. This gene is one of four genes in man and mouse that encode different isoforms of the a subunit. Alternatively spliced transcript variants encoding the same protein have been described. Mutations in this gene are associated with renal tubular acidosis associated with preserved hearing.

## Research Area


## Image Data




Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)



Flow cytometric analysis of HeLa cells using ATP6V0A4 mouse mAb (green) and negative control (red).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissues using ATP6V0A4 mouse mAb with DAB staining.



Immunohistochemical analysis of paraffin-embedded human kidney tissues using ATP6V0A4 mouse mAb with DAB staining.