

# **Product Name: DCLK2 Mouse Monoclonal Antibody**

Catalog #: AMM82895

For research use only.

### **Summary**

**Description** Mouse monoclonal Antibody

**Host** Mouse

**Application** IHC,ELISA,FC

**Reactivity** Human

ConjugationUnconjugatedModificationUnmodifiedIsotypeMouse IgG2aClonalityMonoclonalFormLiquid

**Concentration** 1mg/ml

**Storage** Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

**Buffer** Purified antibody in PBS with 0.05% sodium azide

**Purification** Affinity Purification

# **Application**

**Dilution Ratio** IHC 1:200-1:1000,ELISA 1:5000-1:20000,FC 1:200-1:400

Molecular Weight 83.6kDa

# **Antigen Information**

Gene Name DCLK2

Alternative Names CL2; DCK2; CLIK2; DCDC3; CLICK2; DCDC3B; DCAMKL2; CLICK-II

 Gene ID
 166614.0

 SwissProt ID
 Q8N568

**Immunogen** Purified recombinant fragment of human DCLK2 (AA: 652-766) expressed in mammalian.

#### **Background**


This gene encodes a member of the protein kinase superfamily and the doublecortin family. The protein encoded by this gene contains two N-terminal doublecortin domains, which bind microtubules and regulate microtubule polymerization, a C-terminal serine/threonine protein kinase domain, which shows substantial homology to Ca2+/calmodulin-dependent protein




kinase, and a serine/proline-rich domain in between the doublecortin and the protein kinase domains, which mediates multiple protein-protein interactions. The microtubule-polymerizing activity of the encoded protein is independent of its protein kinase activity. Mouse studies show that the DCX gene, another family member, and this gene share function in the establishment of hippocampal organization and that their absence results in a severe epileptic phenotype and lethality, as described in human patients with lissencephaly. Multiple alternatively spliced transcript variants have been identified.

#### **Research Area**


#### **Image Data**



Black line: Control Antigen (100 ng);Purple line: Antigen (10ng); Blue line: Antigen (50 ng); Red line:Antigen (100 ng)



Flow cytometric analysis of U937 cells using DCLK2 mouse mAb (green) and negative control (red).



Immunohistochemical analysis of paraffin-embedded human brain tissues using DCLK2 mouse mAb with DAB staining.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838