

Citrate Antigen Retrieval Buffer (10×, pH 6.0)

Catalog No.: PA10019

Basic Information

Product name	Citrate Antigen Retrieval Buffer (10×, pH 6.0)
Sizes	100 mL, 500 mL
Storage	RT
Shipping	RT
Validity	12 months

Product Introduction

After fixation with paraformaldehyde, formalin or other aldehyde-based fixatives, proteins become cross-linked and carboxymethylated, masking epitopes and causing false-negative or weak signals in immunostaining. These cross-links can be reversed under high temperature or enzymatic conditions, allowing proteins to regain their native conformation. This process—antigen retrieval (AR)—is essential for reliable IHC/IF results. Heat-induced epitope retrieval (HIER) or enzymatic retrieval can be selected according to tissue and antigen characteristics, and the retrieval buffer must be at an appropriate pH.

This product is a 10× citrate antigen retrieval buffer (pH 6.0). Dilute 1:10 to obtain the working solution, suitable for heat-mediated antigen retrieval.

Working Solution Preparation

Add 10 mL of the 10× concentrate to 90 mL distilled water and mix well.

Notes

1. This product is a concentrate and must be diluted before use. The diluted working solution should be used the same day; short-term storage at 4 ° C is acceptable.
2. Select the antigen retrieval buffer that matches the antigen and tissue characteristics.
3. Wear a lab coat and disposable gloves for your safety and health.
4. For research use only; not intended for clinical diagnosis or therapy.

This product is for research use only!