
Product Name: HuB/HuD Rabbit Polyclonal Antibody**Catalog #: APRab00382**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% sodium azide, pH 7.3.
Purification	Affinity Chromatography

Application

Dilution Ratio	WB 1:500-1:1000,IHC 1:50-1:100,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
Molecular Weight	Calculated MW: 40 kDa; Observed MW: 40 kDa

Antigen Information

Gene Name	ELAVL2/ELAVL4
Alternative Names	ELAVL2; HUB; ELAV-like protein 2; ELAV-like neuronal protein 1; Hu-antigen B; HuB; Nervous system-specific RNA-binding protein Hel-N1; ELAVL4; HUD; PNEM; ELAV-like protein 4; Hu-antigen D; HuD; Paraneoplastic encephalomyelitis antigen HuD
Gene ID	1993/1996
SwissProt ID	Q12926/P26378
Immunogen	The antiserum was produced against synthesized peptide derived from human ELAV2/4. AA range:61-110


Background

Eukaryotic translation initiation factor-5 (EIF5) interacts with the 40S initiation complex to promote hydrolysis of bound GTP with concomitant joining of the 60S ribosomal subunit to the 40S initiation complex. The resulting functional 80S ribosomal initiation complex is then active in peptidyl transfer and chain elongations (summary by Si et al. 1996 [PubMed 8663286]).


Research Area

Neuroscience


Image Data




Western blot analysis of HuB/HuD in HeLa lysates using HuB/HuD antibody.



Immunofluorescence analysis of HuB/HuD in HepG2 using ELAV2/4 antibody. The picture on the right is blocked using the synthesized peptide.



Western blot analysis of HuB/HuD in Jurkat and HUVEC lysates using ELAV2/4 antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of HuB/HuD in 293 lysates using HuB/HuD antibody