Product Name: LC3A/B Rabbit Polyclonal Antibody

Catalog #: APRab03870



Summary

Production Name LC3A/B Rabbit Polyclonal Antibody

Description Rabbit polyclonal Antibody

Host Rabbit
Application WB

Reactivity Human, Mouse, Rat

Performance

ConjugationUnconjugatedModificationUnmodified

Isotype IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% sodium **Buffer**

azide, pH 7.3.

Purification Affinity Purification

Immunogen

Gene Name MAP1LC3A/MAP1LC3B

LC3; LC3A; ATG8E; MAP1ALC3; MAP1BLC3; MAP1LC3A; LC3B; ATG8F; MAP1LC3B-a; Alternative Names

MAP1A/1BLC3; MAP1LC3B

Gene ID 84557/81631

SwissProt ID Q9H492/Q9GZQ8.

Application

Dilution Ratio WB 1:500-1:1000

Molecular Weight Calculated MW: 14 kDa; Observed MW: 14,16 kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: LC3A/B Rabbit Polyclonal Antibody

Catalog #: APRab03870



Background


Macroautophagy is the major inducible pathway for the general turnover of cytoplasmic constituents in eukaryotic cells, it is also responsible for the degradation of active cytoplasmic enzymes and organelles during nutrient starvation.

Macroautophagy involves the formation of double-membrane bound autophagosomes which enclose the cytoplasmic constituent targeted for degradation in a membrane bound structure, which then fuse with the lysosome (or vacuole) releasing a single-membrane bound autophagic bodies which are then degraded within the lysosome (or vacuole). MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. These proteins are involved in formation of autophagosomal vacuoles (autophagosomes). MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. MAP1LC3a is one of the light chain subunits and can associate with either MAP1A or MAP1B. The precursor molecule is cleaved by APG4B/ATG4B to form the cytosolic form, LC3-I. This is activated by APG7L/ATG7, transferred to ATG3 and conjugated to phospholipid to form the membrane-bound form, LC3-II.


Research Area

Signal Transduction

Image Data



Western blot analysis of LC3A/B in rat Brain, C2C12, 3T3 and Hela lysates using LC3A/B antibody.




Western blot analysis of LC3A/B in 3T3 and rat Brain lysates using LC3A/B antibody.

Product Name: LC3A/B Rabbit Polyclonal Antibody

Catalog #: APRab03870





Western blot analysis of LC3A/B in rat Hippocampus, rat Cerebral Cortex and rat Brain lysates using LC3A/B antibody.

Note

For research use only.