# Product Name: Cdc25C (phospho Ser198) Rabbit

Polyclonal Antibody Catalog #: APRab04423



#### **Summary**

Production Name Cdc25C (phospho Ser198) Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

**Host** Rabbit

**Application** IHC-P,IF-P,IF-F,ICC/IF,ELISA

**Reactivity** Human, Rat, Mouse

#### **Performance**

ConjugationUnconjugatedModificationPhosphorylated

**Isotype** IgG

Clonality Polyclonal Form Liquid

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type

preservative N.

**Purification** Affinity purification

#### **Immunogen**

**Buffer** 

Gene Name CDC25C

Alternative Names CDC25C; M-phase inducer phosphatase 3; Dual specificity phosphatase Cdc25C

**Gene ID** 995.0

P30307. The antiserum was produced against synthesized peptide derived from human **SwissProt ID** 

CDC25C around the phosphorylation site of Ser198. AA range:164-213

### **Application**

**Dilution Ratio** IHC-P 1:100-1:300, ELISA 1:5000, IF-P/IF-F/ICC/IF 1:50-200

**Molecular Weight** 

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Cdc25C (phospho Ser198) Rabbit

Polyclonal Antibody Catalog #: APRab04423




## **Background**

cell division cycle 25C(CDC25C) Homo sapiens This gene encodes a conserved protein that plays a key role in the regulation of cell division. The encoded protein directs dephosphorylation of cyclin B-bound CDC2 and triggers entry into mitosis. It also suppresses p53-induced growth arrest. Multiple alternatively spliced transcript variants of this gene have been described. [provided by RefSeq, Dec 2015],catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate.,developmental stage:Expressed predominantly in G2 phase.,function:Functions as a dosage-dependent inducer in mitotic control. It is a tyrosine protein phosphatase required for progression of the cell cycle. It directly dephosphorylates CDC2 and activate its kinase activity.,PTM:Phosphorylated by CHK1 on Ser-216. This phosphorylation creates a binding site for 14-3-3 protein and inhibits the phosphatase.,similarity:Belongs to the MPI phosphatase family,,similarity:Contains 1 rhodanese domain,,subunit:Interacts with HIV-1 Vpr, thereby inactivating CDC25C phosphatase activity.,

#### Research Area

Cell\_Cycle\_G1S;Cell\_Cycle\_G2M\_DNA;Oocyte meiosis;Progesterone-mediated oocyte maturation;

## **Image Data**



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using CDC25C (Phospho-Ser198)

Antibody. The picture on the right is blocked with the phospho peptide.

#### **Note**

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838