

# Product Name: Ephrin-B1 (phospho Tyr317) Rabbit Polyclonal Antibody Catalog #: APRab04617

For research use only.

#### **Summary**

**Description** Rabbit polyclonal Antibody

Host Rabbit
Application WB,ELISA

Reactivity Human, Mouse, Rat
Conjugation Unconjugated
Modification Phosphorylated

**Isotype** IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

**Storage** Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

#### **Application**

**Dilution Ratio** WB 1:500-1:2000,ELISA 1:10000-1:20000

Molecular Weight 30kDa

### **Antigen Information**

**Alternative Names** 

Gene Name EFNB1

EFNB1; EFL3; EPLG2; LERK2; Ephrin-B1; EFL-3; ELK ligand; ELK-L; EPH-related receptor

tyrosine kinase ligand 2; LERK-2

 Gene ID
 1947.0

 SwissProt ID
 P98172

The antiserum was produced against synthesized peptide derived from human EFNB1 Immunogen

around the phosphorylation site of Tyr317. AA range:283-332

## **Background**




The protein encoded by this gene is a type I membrane protein and a ligand of Eph-related receptor tyrosine kinases. It may play a role in cell adhesion and function in the development or maintenance of the nervous system. [provided by RefSeq, Jul 2008], disease: Defects in EFNB1 are a cause of craniofrontonasal syndrome (CFNS) [MIM:304110]; also known as craniofrontonasal dysplasia (CFND). CFNS is an X-linked inherited syndrome characterized by hypertelorism, coronal synostosis with brachycephaly, downslanting palpebral fissures, clefting of the nasal tip, joint anomalies, longitudinally grooved fingernails and other digital anomalies., function: Binds to the receptor tyrosine kinases EPHB1 and EPHA1. Binds to, and induce the collapse of, commissural axons/growth cones in vitro. May play a role in constraining the orientation of longitudinally projecting axons., induction: By TNF-alpha., PTM: Inducible phosphorylation of tyrosine residues in the cytoplasmic domain., similarity: Belongs to the ephrin family., subunit: Interacts with GRIP1 and GRIP2., tissue specificity: Heart, placenta, lung, liver, skeletal muscle, kidney, pancreas.,


#### **Research Area**

Axon guidance;

#### **Image Data**




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using EFNB1 (Phospho-Tyr317) Antibody



Western blot analysis of lysates from mouse brain, using EFNB1 (Phospho-Tyr317) Antibody. The lane on the right is blocked with the phospho peptide.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Western Blot analysis of mouse cells using Phospho-Ephrin-B1 (Y317) Polyclonal Antibody