

Product Name: FGFR-4 (phospho Tyr642) Rabbit Polyclonal Antibody**Catalog #: APRab04670**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Phosphorylated
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight	90kDa

Antigen Information

Gene Name	FGFR4
Alternative Names	FGFR4; JTK2; TKF; Fibroblast growth factor receptor 4; FGFR-4; CD antigen CD334
Gene ID	2264.0
SwissProt ID	P22455
Immunogen	Synthesized phospho-peptide around the phosphorylation site of human FGFR-4 (phospho Tyr642)

Background


The protein encoded by this gene is a member of the fibroblast growth factor receptor family, where amino acid sequence is

highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. The genomic organization of this gene, compared to members 1-3, encompasses 18 exons rather than 19 or 20. Although alternative splicing has been observed, there is no evidence that the C-terminal half of the IgII catalytic activity: $\text{ATP} + \text{a [protein]-L-tyrosine} = \text{ADP} + \text{a [protein]-L-tyrosine phosphate}$. function: Receptor for acidic fibroblast growth factor. Does not bind to basic fibroblast growth factor. Binds FGF19. PTM: Glycosylated (By similarity). Phosphorylated on tyrosine residue (By similarity). Phosphorylation requires the presence of a functional (phosphorylated) FGFR1 and not necessarily by means of FGFR heterodimerization. similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. Fibroblast growth factor receptor subfamily. similarity: Contains 1 protein kinase domain. similarity: Contains 3 Ig-like C2-type (immunoglobulin-like) domains. subcellular location: Isoform 2 may be secreted. subunit: Interacts with KLB. tissue specificity: Expressed in gastrointestinal epithelial cells, pancreas, and gastric and pancreatic cancer cell lines.

Research Area

MAPK_ERK_Growth; MAPK_G_Protein; Endocytosis; Regulates Actin and Cytoskeleton;

Image Data



Western Blot analysis of JK cells using Phospho-FGFR-4 (Y642) Polyclonal Antibody