Polyclonal Antibody Catalog #: APRab04704



Summary

Production Name FoxO4 (phospho Ser197) Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA

Reactivity Human, Mouse

Performance

ConjugationUnconjugatedModificationPhosphorylated

Isotype IgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type

preservative N.

Purification Affinity purification

Immunogen

Gene Name FOXO4

FOXO4; AFX; AFX1; MLLT7; Forkhead box protein O4; Fork head domain transcription

factor AFX1

Gene ID 4303.0

P98177.The antiserum was produced against synthesized peptide derived from human **SwissProt ID**

AFX around the phosphorylation site of Ser197. AA range:164-213

Application

Dilution Ratio WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:20000, IF-P/IF-F/ICC/IF 1:50-200

Molecular Weight 70kDa

Polyclonal Antibody Catalog #: APRab04704



Background

This gene encodes a member of the O class of winged helix/forkhead transcription factor family. Proteins encoded by this class are regulated by factors involved in growth and differentiation indicating they play a role in these processes. A translocation involving this gene on chromosome X and the homolog of the Drosophila trithorax gene, encoding a DNA binding protein, located on chromosome 11 is associated with leukemia. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2010], disease: A chromosomal aberration involving FOXO4 is found in acute leukemias. Translocation t(X;11)(q13;q23) with MLL/HRX. The result is a roque activator protein, function: Transcription factor involved in the regulation of the insulin signaling pathway. Binds to insulin-response elements (IREs) and can activate transcription of IGFBP1. Down-regulates expression of HIF1A and suppresses hypoxiainduced transcriptional activation of HIF1A-modulated genes. Also involved in negative regulation of the cell cycle., pharmaceutical: A constitutively active FOXO4 mutant where phosphorylation sites Thr-32, Ser-187 and Ser-262 have been mutated to alanine may have therapeutic potential in ERBB2/HER2-overexpressing cancers as it inhibits ERBB2mediated cell survival, transformation and tumorigenicity.,PTM:Acetylation by CBP, which is induced by peroxidase stress, inhibits transcriptional activity. Deacetylation by SIRT1 is NAD-dependent and stimulates transcriptional activity, PTM: Phosphorylation by PKB/AKT1 inhibits transcriptional activity and is responsible for cytoplasmic localization., similarity: Contains 1 fork-head DNA-binding domain., subcellular location: When phosphorylated, translocated from nucleus to cytoplasm. Dephosphorylation triggers nuclear translocation., subunit: Interacts with CBP, MYOCD, SIRT1, SRF and YWHAZ. Acetylated by CBP and deacetylated by SIRT1. Binding of YWHAZ inhibits DNA-binding, tissue specificity: Heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Isoform zeta is most abundant in the liver, kidney, and pancreas.,

Research Area


Insulin Receptor; B Cell Receptor; Protein_Acetylation

Image Data


Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Polyclonal Antibody Catalog #: APRab04704






Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using AFX (Phospho-Ser197) Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using AFX (Phospho-Ser197) Antibody.


The picture on the right is blocked with the phospho peptide.




Western blot analysis of lysates from 293 cells treated with serum, using AFX (Phospho-Ser197) Antibody. The lane on the right is blocked with the phospho peptide.

Polyclonal Antibody Catalog #: APRab04704





Western Blot analysis of various cells using Phospho-FoxO4 (S197) Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

Note

For research use only.