Product Name: p70 S6 kinase α (phospho Ser434) Rabbit Polyclonal Antibody

Polyclonal Antibody Catalog #: APRab05188



Summary

Production Name p70 S6 kinase α (phospho Ser434) Rabbit Polyclonal Antibody

Description Rabbit Polyclonal Antibody

Host Rabbit

Application WB,IHC-P,IF-P,IF-F,ICC/IF,ELISA

Reactivity Human, Mouse, Rat, Monkey

Performance

ConjugationUnconjugatedModificationPhosphorylated

Isotype IgG

Clonality Polyclonal Form Liquid

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type

preservative N.

Purification Affinity purification

Immunogen

Buffer

Gene Name RPS6KB1

RPS6KB1; STK14A; Ribosomal protein S6 kinase beta-1; S6K-beta-1; S6K1; 70 kDa

Alternative Names ribosomal protein S6 kinase 1; P70S6K1; p70-S6K 1; Ribosomal protein S6 kinase I;

Serine/threonine-protein kinase 14A; p70 ribosomal S6 kinase alpha; p70 S6 kinas

Gene ID 6198.0

P23443. The antiserum was produced against synthesized peptide derived from human

p70 S6 Kinase around the phosphorylation site of Ser434. AA range:401-450

Application

SwissProt ID

Dilution Ratio WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:10000, IF-P/IF-F/ICC/IF 1:50-200

Molecular Weight 60kDa

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: p70 S6 kinase α (phospho Ser434) Rabbit

Polyclonal Antibody Catalog #: APRab05188




Background

ribosomal protein S6 kinase B1(RPS6KB1) Homo sapiens This gene encodes a member of the ribosomal S6 kinase family of serine/threonine kinases. The encoded protein responds to mTOR (mammalian target of rapamycin) signaling to promote protein synthesis, cell growth, and cell proliferation. Activity of this gene has been associated with human cancer. Alternatively spliced transcript variants have been observed. The use of alternative translation start sites results in isoforms with longer or shorter N-termini which may differ in their subcellular localizations. There are two pseudogenes for this gene on chromosome 17. [provided by RefSeq, Jan 2013],catalytic activity:ATP + a protein = ADP + a phosphoprotein,enzyme regulation:Activation by serine/threonine phosphorylation and protein kinase C, inactivated by type 2A phosphatase,function:Phosphorylates specifically ribosomal protein S6 in response to insulin or several classes of mitogens, similarity:Belongs to the protein kinase superfamily, similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. S6 kinase subfamily, similarity:Contains 1 AGC-kinase C-terminal domain, similarity:Contains 1 protein kinase domain, subunit:Interacts with PPP1R9A/neurabin-1, tissue specificity:Widely expressed.

Research Area

Regulates Angiogenesis; Insulin Receptor; ErbB/HER; mTOR; B Cell Receptor; PI3K/Akt; PI3K/Akt; AMPK

Image Data



Western blot analysis of lysates from K562 cells, NIH/3T3 cells and COLO205 cells, using p70 S6 Kinase (Phospho-Ser411)

Antibody. The lane on the right is blocked with the phospho peptide.

Note

For research use only.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838