
Product Name: WWOX (phospho Tyr33) Rabbit Polyclonal Antibody**Catalog #: APRab05635**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Phosphorylated
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
Molecular Weight	55kDa

Antigen Information


Gene Name	WWOX
Alternative Names	WWOX; FOR; WOX1; WW domain-containing oxidoreductase; Fragile site FRA16D oxidoreductase
Gene ID	51741.0
SwissProt ID	Q96KM3
Immunogen	The antiserum was produced against synthesized peptide derived from human WWOX around the phosphorylation site of Tyr33. AA range:18-67

Background


disease:Defects in WWOX may be involved in esophageal squamous cell carcinoma (ESCC) [MIM:133239],disease:Defects in WWOX may be involved in several cancer types. The gene spans the second most common chromosomal fragile site (FRA16D) which is frequently altered in cancers. Alteration of the expression and expression of some isoforms is associated with cancers. However, it is still unclear if alteration of WWOX is directly implicated in cancerogenesis or if it corresponds to a secondary effect.,domain:The WW 1 domain mediates interaction with TP53, and probably TP73, TFAP2C, LITAF and WBP1.,function:Probable oxidoreductase, which acts as a tumor suppressor and plays a role in apoptosis. May function synergistically with TP53/p53 to control genotoxic stress-induced cell death. May also play a role in tumor necrosis factor (TNF)-mediated cell death.,PTM:Phosphorylated upon genotoxic stress. Phosphorylation of Tyr-33 regulates interaction with TP53, TP73 and MAPK8. May also regulate proapoptotic activity.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,similarity:Contains 2 WW domains.,subcellular location:Partially localizes to the mitochondria. Translocates to the nucleus upon genotoxic stress or TNF stimulation (By similarity). Isoform 5 and isoform 6 may localize in the nucleus.,subunit:Interacts with TP53, TP73/p73 and MAPK8. Interacts with MAPT/TAU (By similarity). Forms a ternary complex with TP53 and MDM2. Interacts with ERBB4, LITAF and WBP1. May interact with COTE1/C1orf2 and SCOTIN.,tissue specificity:Widely expressed. Strongly expressed in testis, prostate, and ovary. Overexpressed in cancer cell lines. Isoform 5 and isoform 6 may only be expressed in tumor cell lines.,disease:Defects in WWOX may be involved in esophageal squamous cell carcinoma (ESCC) [MIM:133239],disease:Defects in WWOX may be involved in several cancer types. The gene spans the second most common chromosomal fragile site (FRA16D) which is frequently altered in cancers. Alteration of the expression and expression of some isoforms is associated with cancers. However, it is still unclear if alteration of WWOX is directly implicated in cancerogenesis or if it corresponds to a secondary effect.,domain:The WW 1 domain mediates interaction with TP53, and probably TP73, TFAP2C, LITAF and WBP1.,function:Probable oxidoreductase, which acts as a tumor suppressor and plays a role in apoptosis. May function synergistically with TP53/p53 to control genotoxic stress-induced cell death. May also play a role in tumor necrosis factor (TNF)-mediated cell death.,PTM:Phosphorylated upon genotoxic stress. Phosphorylation of Tyr-33 regulates interaction with TP53, TP73 and MAPK8. May also regulate proapoptotic activity.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,similarity:Contains 2 WW domains.,subcellular location:Partially localizes to the mitochondria. Translocates to the nucleus upon genotoxic stress or TNF stimulation (By similarity). Isoform 5 and isoform 6 may localize in the nucleus.,subunit:Interacts with TP53, TP73/p73 and MAPK8. Interacts with MAPT/TAU (By similarity). Forms a ternary complex with TP53 and MDM2. Interacts with ERBB4, LITAF and WBP1. May interact with COTE1/C1orf2 and SCOTIN.,tissue specificity:Widely expressed. Strongly expressed in testis, prostate, and ovary. Overexpressed in cancer cell lines. Isoform 5 and isoform 6 may only be expressed in tumor cell lines.,

Research Area

Image Data



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using WWOX (Phospho-Tyr33) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HepG2 cells treated with PMA 125ng/ml 30', using WWOX (Phospho-Tyr33) Antibody. The lane on the right is blocked with the phospho peptide.