

---

**Product Name: 7B2 Rabbit Polyclonal Antibody****Catalog #: APRab06349**

For research use only.

**Summary**

<b>Description</b>	Rabbit polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human,Mouse,Rat
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

**Application**

<b>Dilution Ratio</b>	WB 1:500-1:2000,ELISA 1:5000-1:20000
<b>Molecular Weight</b>	23kDa

**Antigen Information**

<b>Gene Name</b>	SCG5 SGNE1
<b>Alternative Names</b>	
<b>Gene ID</b>	6447.0
<b>SwissProt ID</b>	P05408
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 90-170


**Background**

This gene encodes a secreted chaperone protein that prevents the aggregation of other secreted proteins, including proteins that are associated with neurodegenerative and metabolic disease. The encoded protein may be best known for its role in the trafficking and activation of prohormone convertase PC2 (encoded by Gene ID: 5126). Phosphorylation of the encoded protein

has been shown to have an inhibitory effect on its chaperone function. [provided by RefSeq, Jul 2016],function:Acts as a molecular chaperone for PCSK2/PC2, preventing its premature activation in the regulated secretory pathway. Binds to inactive PCSK2 in the endoplasmic reticulum and facilitates its transport from there to later compartments of the secretory pathway where it is proteolytically matured and activated. Also required for cleavage of PCSK2 but does not appear to be involved in its folding. Plays a role in regulating pituitary hormone secretion. The C-terminal peptide inhibits PCSK2 in vitro.,PTM:Proteolytically cleaved in the Golgi by a furin-like convertase to generate bioactive peptides.,PTM:Sulfated on tyrosine residues.,similarity:Belongs to the 7B2 family.,subcellular location:Neuroendocrine and endocrine secretory granules.,subunit:Interacts with PCSK2/PC2 early in the secretory pathway. Dissociation occurs at later stages.,

## Research Area

## Image Data



Western blot analysis of lysates from K562 cells, 7B2 Rabbit Polyclonal Antibody was diluted at 1:1000, 4° over night