
Product Name: 9G8 Rabbit Polyclonal Antibody**Catalog #: APRab06351**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
Molecular Weight	35kDa

Antigen Information

Gene Name	SRSF7
Alternative Names	SRSF7; SFRS7; Serine/arginine-rich splicing factor 7; Splicing factor 9G8; Splicing factor; arginine/serine-rich 7
Gene ID	6432.0
SwissProt ID	Q16629
Immunogen	The antiserum was produced against synthesized peptide derived from human SFRS7. AA range:61-110


Background

The protein encoded by this gene is a member of the serine/arginine (SR)-rich family of pre-mRNA splicing factors, which constitute part of the spliceosome. Each of these factors contains an RNA recognition motif (RRM) for binding RNA and an RS domain for binding other proteins. The RS domain is rich in serine and arginine residues and facilitates interaction between different SR splicing factors. In addition to being critical for mRNA splicing, the SR proteins have also been shown to be involved in mRNA export from the nucleus and in translation. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2010], alternative products: Isoforms, often lacking the RS domain and differentially expressed in fetal tissues, may be involved in modulation of 9G8 function, function: Required for pre-mRNA splicing. Can also modulate alternative splicing in vitro., PTM: Extensively phosphorylated on serine residues in the RS domain., similarity: Belongs to the splicing factor SR family., similarity: Contains 1 CCHC-type zinc finger., similarity: Contains 1 RRM (RNA recognition motif) domain., subunit: Found in large molecular weight complexes containing CCNL1 and the p110 isoforms of either CDC2L1 or CDC2L2. Interacts with CCNL2 and CPSF6., tissue specificity: Brain, liver, kidney and lung.,


Research Area

Spliceosome;

Image Data




Western blot analysis of lysates from HT-29 cells, using SFRS7 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using SFRS7 antibody.

Western Blot analysis of various cells using 9G8 Polyclonal Antibody.



Western Blot analysis of HuvEc cells using 9G8 Polyclonal Antibody.

