

Product Name: ACAD-11 Rabbit Polyclonal Antibody**Catalog #: APRab06459**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight	87kDa

Antigen Information

Gene Name	ACAD11
Alternative Names	ACAD11; Acyl-CoA dehydrogenase family member 11; ACAD-11
Gene ID	84129.0
SwissProt ID	Q709F0
Immunogen	The antiserum was produced against synthesized peptide derived from human ACAD11. AA range:381-430


Background

acyl-CoA dehydrogenase family member 11(ACAD11) Homo sapiens This gene encodes an acyl-CoA dehydrogenase enzyme


with a preference for carbon chain lengths between 20 and 26. Naturally occurring read-through transcription occurs between the upstream gene NPHP3 (nephronophthisis 3 (adolescent)) and this gene. [provided by RefSeq, Aug 2015],alternative products:Additional isoforms seem to exist,disease:Defects in NPHP3 are a cause of renal-hepatic-pancreatic dysplasia (RHPD) [MIM:208540]. RHPD is an autosomal recessive disorder with variable expression, and patients surviving the neonatal period progress to renal and hepatic failure which can be treated successfully with combined liver-kidney transplantation.,disease:Defects in NPHP3 are the cause of nephronophthisis type 3 (NPHP3) [MIM:604387]; also known as adolescent nephronophthisis. NPHP3 is a autosomal recessive disorder resulting in end-stage renal disease. It is characterized by polyuria, polydipsia, anemia. Onset of terminal renal failure occurs significantly later (median age, 19 years) than in juvenile nephronophthisis. Renal pathology is characterized by alterations of tubular basement membranes, tubular atrophy and dilatation, sclerosing tubulointerstitial nephropathy, and renal cyst development predominantly at the corticomedullary junction.,function:May participate in mechanosensation in the primary cilium of kidney cells.,similarity:Belongs to the acyl-CoA dehydrogenase family.,similarity:Contains 11 TPR repeats.,subunit:Interacts with NPHP1.,tissue specificity:Widely expressed at low level. Expressed in heart, placenta, liver, skeletal muscle, kidney and pancreas. Expressed at very low level in brain and lung.,

Research Area

Image Data



Western blot analysis of ACAD11 Antibody. The lane on the right is blocked with the ACAD11 peptide.



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using ACAD11 Antibody. The lane on the right is blocked with the ACAD11 peptide.