
Product Name: ACAD-9 Rabbit Polyclonal Antibody**Catalog #: APRab06460**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	IHC,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000

Molecular Weight

Antigen Information

Gene Name	ACAD9
Alternative Names	ACAD9; Acyl-CoA dehydrogenase family member 9; mitochondrial; ACAD-9
Gene ID	28976.0
SwissProt ID	Q9H845
Immunogen	Synthesized peptide derived from ACAD-9 . at AA range: 530-610


Background

This gene encodes a member of the acyl-CoA dehydrogenase family. Members of this family of proteins localize to the mitochondria and catalyze the rate-limiting step in the beta-oxidation of fatty acyl-CoA. The encoded protein is specifically

active toward palmitoyl-CoA and long-chain unsaturated substrates. Mutations in this gene cause acyl-CoA dehydrogenase family member type 9 deficiency. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Mar 2010],cofactor:FAD.,disease:Defects in ACAD9 are a cause of acyl-CoA dehydrogenase family member type 9 deficiency (ACAD9 deficiency) [MIM:611126]. ACAD9 deficiency patients present with episodic liver dysfunction during otherwise mild illnesses or cardiomyopathy, along with chronic neurologic dysfunction.,function:Has a dehydrogenase activity on palmitoyl-CoA (C16:0) and stearoyl-CoA (C18:0). It is three times more active on palmitoyl-CoA than on stearoyl-CoA. Has little activity on octanoyl-CoA (C8:0), butyryl-CoA (C4:0) or isovaleryl-CoA (5:0).,similarity:Belongs to the acyl-CoA dehydrogenase family.,tissue specificity:Ubiquitously expressed in most normal human tissues and cancer cell lines with high level of expression in heart, skeletal muscles, brain, kidney and liver.,

Research Area

Image Data



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .