
Product Name: ADAMTS-12 Rabbit Polyclonal Antibody**Catalog #: APRab06596**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	IHC,ICC/IF,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000

Molecular Weight

Antigen Information

Gene Name	ADAMTS12
Alternative Names	ADAMTS12; A disintegrin and metalloproteinase with thrombospondin motifs 12; ADAM-TS 12; ADAM-TS12; ADAMTS-12
Gene ID	81792.0
SwissProt ID	P58397
Immunogen	Synthesized peptide derived from ADAMTS-12 . at AA range: 1100-1180


Background

This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motifs) protein family.

Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS-1) motif. Individual members of this family differ in the number of C-terminal TS-1 motifs, and some have unique C-terminal domains. The enzyme encoded by this gene contains eight TS-1 motifs. It may play roles in pulmonary cells during fetal development or in tumor processes through its proteolytic activity or as a molecule potentially involved in regulation of cell adhesion. [provided by RefSeq, Jul 2008],cofactor: Binds 1 zinc ion per subunit.,domain: The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,domain: The spacer domain and the TSP type-1 domains are important for a tight interaction with the extracellular matrix.,PTM: Subjected to an intracellular maturation process yielding a 120 kDa N-terminal fragment containing the metalloproteinase, disintegrin, one TSP type-1 and the Cys-rich domains and a 83 kDa C-terminal fragment containing the spacer 2 and four TSP type-1 domains.,PTM: The precursor is cleaved by a furin endopeptidase.,similarity: Contains 1 disintegrin domain.,similarity: Contains 1 peptidase M12B domain.,similarity: Contains 1 PLAC domain.,similarity: Contains 8 TSP type-1 domains.,tissue specificity: Expressed exclusively in fetal lung. Is widely expressed in gastric carcinomas and in cancer cells of diverse origin.,

Research Area

Image Data



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .