

Product Name: ADAR2 Rabbit Polyclonal Antibody**Catalog #: APRab06605**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight	80kDa

Antigen Information


Gene Name	ADARB1
Alternative Names	ADARB1; ADAR2; DRADA2; RED1; Double-stranded RNA-specific editase 1; RNA-editing deaminase 1; RNA-editing enzyme 1; dsRNA adenosine deaminase
Gene ID	104.0
SwissProt ID	P78563
Immunogen	The antiserum was produced against synthesized peptide derived from human ADARB1. AA range:481-530

Background

This gene encodes the enzyme responsible for pre-mRNA editing of the glutamate receptor subunit B by site-specific deamination of adenosines. Studies in rat found that this enzyme acted on its own pre-mRNA molecules to convert an AA dinucleotide to an AI dinucleotide which resulted in a new splice site. Alternative splicing of this gene results in several transcript variants, some of which have been characterized by the presence or absence of an ALU cassette insert and a short or long C-terminal region. [provided by RefSeq, Jul 2008],alternative products:Additional isoforms seem to exist,cofactor:Binds 1 inositol hexakisphosphate (IP6) per subunit.,function:Editing of the messenger RNAs for glutamate receptor (GluR) subunits by site-selective adenosine deamination. Edits both the GluR-B Q/R and R/G sites efficiently but converts the adenosine in hotspot1 much less efficiently.,similarity:Contains 1 A to I editase domain.,similarity:Contains 2 DRBM (double-stranded RNA-binding) domains.,

Research Area

Image Data



Western blot analysis of lysates from HepG2 cells, using ADARB1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using ADAR2 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA) .