

Product Name: Adenosine A2A-R Rabbit Polyclonal Antibody

Catalog #: APRab06622

For research use only.

Modification

Summary

Description Rabbit polyclonal Antibody

Host Rabbit

ApplicationWB,IHC,ICC/IF,ELISAReactivityHuman,Rat,MouseConjugationUnconjugated

Unmodified

Isotype IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

Shipping Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer**

preservative N.

Purification Affinity purification

Application

Dilution Ratio WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000

Molecular Weight 37kDa

Antigen Information

Gene Name ADORA2A

Alternative Names ADORA2A; ADORA2; Adenosine receptor A2a

 Gene ID
 135.0

 SwissProt ID
 P29274

The antiserum was produced against synthesized peptide derived from human ADORA2A.

AA range:120-169

Background

adenosine A2a receptor(ADORA2A) Homo sapiens This gene encodes a member of the guanine nucleotide-binding protein

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




(G protein)-coupled receptor (GPCR) superfamily, which is subdivided into classes and subtypes. The receptors are seven-pass transmembrane proteins that respond to extracellular cues and activate intracellular signal transduction pathways. This protein, an adenosine receptor of A2A subtype, uses adenosine as the preferred endogenous agonist and preferentially interacts with the G(s) and G(olf) family of G proteins to increase intracellular cAMP levels. It plays an important role in many biological functions, such as cardiac rhythm and circulation, cerebral and renal blood flow, immune function, pain regulation, and sleep. It has been implicated in pathophysiological conditions such as inflammatory diseases and neurodegenerative disorders. Alternative splicing results in multiple transcript variants. A read-through transcript composfunction:Receptor for adenosine. The activity of this receptor is mediated by G proteins which activate adenylyl cyclase.,similarity:Belongs to the G-protein coupled receptor 1 family.,


Research Area

Calcium; Neuroactive ligand-receptor interaction; Vascular smooth muscle contraction;


Image Data




Immunofluorescence analysis of COS7 cells, using ADORA2A Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using ADORA2A Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of ADORA2A Antibody. The lane on the right is blocked with the ADORA2A peptide.




Immunohistochemistryt analysis of paraffin-embedded human brain, using ADORA2A Antibody. The lane on the right is blocked with the ADORA2A peptide.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Immunofluorescence analysis of ADORA2A Antibody. The lane on the right is blocked with the ADORA2A peptide.



Western blot analysis of the lysates from HepG2 cells using ADORA2A antibody.