

Product Name: AKR1CL2 Rabbit Polyclonal Antibody

Catalog #: APRab06736

For research use only.

Summary

Description Rabbit polyclonal Antibody

Host Rabbit

ApplicationWB,IHC,ICC/IF,ELISAReactivityHuman,Rat,MouseConjugationUnconjugatedModificationUnmodified

Isotype IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

Shipping Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer**

preservative N.

Purification Affinity purification

Application

Dilution Ratio WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000

Molecular Weight 37kDa

Antigen Information

Gene Name AKR1E2

AKR1E2; AKR1CL2; AKRDC1; 1; 5-anhydro-D-fructose reductase; AF reductase; Aldo-keto

Alternative Names reductase family 1 member C-like protein 2; Aldo-keto reductase family 1 member E2;

LoopADR; Testis-specific protein; hTSP

 Gene ID
 83592.0

 SwissProt ID
 Q96JD6

The antiserum was produced against synthesized peptide derived from human AKR1CL2. AA

Immunogen range:141-190




Background


The protein encoded by this gene is a member of the aldo-keto reductase superfamily. Members in this family are characterized by their structure (evolutionarily highly conserved TIM barrel) and function (NAD(P)H-dependent oxido-reduction of carbonyl groups). Transcripts of this gene have been reported in specimens of human testis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2012],catalytic activity:1,5-anhydro-D-glucitol + NADP(+) = 1,5-anhydro-D-fructose + NADPH.,function:Catalyzes the NADPH-dependent reduction of 1,5-anhydro-D-fructose (AF) to 1,5-anhydro-D-glucitol. Can also catalyze the reduction of various aldehydes and quinones.,similarity:Belongs to the aldo/keto reductase family.,tissue specificity:Testis-specific.,

Research Area

Image Data




Western blot analysis of lysates from HepG2 and Jurkat cells, using AKR1CL2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using AKR1CL2 antibody.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Western Blot analysis of various cells using AKR1CL2 Polyclonal Antibody diluted at $1\colon 500$

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838