

Product Name: AMY-1 Rabbit Polyclonal Antibody**Catalog #: APRab06862**

For research use only.

Summary

| | |
|----------------------|---|
| Description | Rabbit polyclonal Antibody |
| Host | Rabbit |
| Application | IHC, ICC/IF, ELISA |
| Reactivity | Human, Mouse |
| Conjugation | Unconjugated |
| Modification | Unmodified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Concentration | 1mg/ml |
| Storage | Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles. |
| Shipping | Ice bags |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| Purification | Affinity purification |

Application

Dilution Ratio IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:5000-1:20000

Molecular Weight

Antigen Information

| | |
|--------------------------|---|
| Gene Name | MYCBP |
| Alternative Names | MYCBP; AMY1; C-Myc-binding protein; Associate of Myc 1; AMY-1 |
| Gene ID | 26292.0 |
| SwissProt ID | Q99417 |
| Immunogen | Synthesized peptide derived from AMY-1 . at AA range: 30-110 |


Background

The protein encoded by this gene binds to the N-terminus of the oncogenic protein C-MYC, enhancing the ability of C-MYC to activate E box-dependent transcription. The encoded protein is normally found in the cytoplasm, but it translocates to the

nucleus during S phase of the cell cycle and associates with C-MYC. This protein may be involved in spermatogenesis. This gene can be silenced by microRNA-22. Two transcript variants, one protein-coding and the other probably not protein-coding, have been found for this gene. [provided by RefSeq, Nov 2011],function:May control the transcriptional activity of MYC. Stimulates the activation of E box-dependent transcription by MYC.,similarity:Belongs to the AMY1 family.,subcellular location:Translocates into the nucleus in the S phase of the cell cycle upon an increase of c-Myc expression. Found in the mitochondria when associated with AKAP1.,subunit:Binds via its C-terminal region to the N-terminal region of MYC. Associates with AKAP1/S-AKAP84. Interacts with MYCBPAP.,tissue specificity:Highly expressed in heart, placenta, pancreas, skeletal muscle and kidney. Also present at low levels in lung,.

Research Area

Image Data



Immunohistochemical analysis of paraffin-embedded human oophoroma. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .