
Product Name: ARHG4 Rabbit Polyclonal Antibody**Catalog #: APRab07116**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight	75kDa

Antigen Information

Gene Name	ARHGEF4 KIAA1112
Alternative Names	
Gene ID	50649.0
SwissProt ID	Q9NR80
Immunogen	Synthesized peptide derived from human protein . at AA range: 270-350

Background


Rho GTPases play a fundamental role in numerous cellular processes that are initiated by extracellular stimuli that work through G protein coupled receptors. The protein encoded by this gene may form complex with G proteins and stimulate Rho-dependent signals. Multiple alternatively spliced transcript variants encoding different isoforms have been found, but the full-

length nature of some variants has not been determined. [provided by RefSeq, Jun 2013],function:Acts as guanine nucleotide exchange factor (GEF) for RhoA and RAC1 GTPases. Binding of APC may activate RAC1 GEF activity. The APC-ARHGEF4 complex seems to be involved in cell migration as well as in E-cadherin-mediated cell-cell adhesion.,sequence caution:Translation N-terminally extended.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 SH3 domain.,subcellular location:Associated with membrane ruffles.,subunit:Isoform 3 interacts with RHOA and RAC1, and through its N-terminus with APC. Found in a complex consisting of ARHGEF4, APC and CTNNB1.,tissue specificity:Expressed at low levels in brain, kidney, lung and muscle.,

Research Area

Regulates Actin and Cytoskeleton;

Image Data



Western Blot analysis of HEK293 lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000