

# **Product Name:** Arrestin-β-1 Rabbit Polyclonal Antibody

Catalog #: APRab07170

For research use only.

### **Summary**

**Description** Rabbit polyclonal Antibody

**Host** Rabbit

Application WB,IHC,ICC/IF,ELISA

Reactivity Human, Monkey
Conjugation Unconjugated
Modification Unmodified

**Isotype** IgG

ClonalityPolyclonalFormLiquidConcentration1mg/ml

Storage Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.

**Shipping** Ice bags

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

# **Application**

**Dilution Ratio** WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000

Molecular Weight 47kDa

# **Antigen Information**

Gene Name ARRB1

Alternative Names ARRB1; ARR1; Beta-arrestin-1; Arrestin beta-1

 Gene ID
 408.0

 SwissProt ID
 P49407

The antiserum was produced against synthesized peptide derived from human Arrestin 1. AA Immunogen

range:369-418

## **Background**

Members of arrestin/beta-arrestin protein family are thought to participate in agonist-mediated desensitization of G-protein-

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838




coupled receptors and cause specific dampening of cellular responses to stimuli such as hormones, neurotransmitters, or sensory signals. Arrestin beta 1 is a cytosolic protein and acts as a cofactor in the beta-adrenergic receptor kinase (BARK) mediated desensitization of beta-adrenergic receptors. Besides the central nervous system, it is expressed at high levels in peripheral blood leukocytes, and thus the BARK/beta-arrestin system is believed to play a major role in regulating receptor-mediated immune functions. Alternatively spliced transcripts encoding different isoforms of arrestin beta 1 have been described. [provided by RefSeq, Jan 2011],function:Regulates beta-adrenergic receptor function. Beta-arrestins seem to bind phosphorylated beta-adrenergic receptors, thereby causing a significant impairment of their capacity to activate G(S) proteins, online information:Arrestin entry, similarity:Belongs to the arrestin family.


#### **Research Area**

MAPK\_ERK\_Growth;MAPK\_G\_Protein;Chemokine;Endocytosis;


#### **Image Data**



Immunofluorescence analysis of A549 cells, using Arrestin 1 Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Arrestin 1 Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from A549 cells, treated with Etoposide 25uM 60  $\,^{'}$ , using Arrestin 1 Antibody. The lane on the right is blocked with the synthesized peptide.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838





Western Blot analysis of various cells using Arrestin- $\beta$ -1 Polyclonal Antibody diluted at 1:  $\,500\,$ 



Western Blot analysis of various cells using Antibody diluted at 1:1000. Secondary antibody was diluted at 1:20000