
Product Name: Ataxin-2L Rabbit Polyclonal Antibody**Catalog #: APRab07253**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:10000-1:20000
Molecular Weight	113kDa

Antigen Information


Gene Name	ATXN2L
Alternative Names	ATXN2L; A2D; A2LG; A2LP; A2RP; Ataxin-2-like protein; Ataxin-2 domain protein; Ataxin-2-related protein
Gene ID	11273.0
SwissProt ID	Q8WWM7
Immunogen	The antiserum was produced against synthesized peptide derived from human ATXN2L. AA range:571-620

Background


This gene encodes an ataxin type 2 related protein of unknown function. This protein is a member of the spinocerebellar ataxia (SCAs) family, which is associated with a complex group of neurodegenerative disorders. Several alternatively spliced transcripts encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],PTM:Arg-361 is dimethylated, probably to asymmetric dimethylarginine.,PTM:Thrombopoietin triggers the phosphorylation on tyrosine residues in a way that is dependent on MPL C-terminal domain.,sequence caution:Exon duplication.,similarity:Belongs to the ataxin-2 family.,subunit:Interacts with MPL/TPOR and EPOR and dissociates after ligand stimulation.,tissue specificity:Expressed at high levels in thymus, lymph node, spleen, fetal kidney and adult testis. Constitutively associated with MPL and EPOR in hematopoietic cells.,

Research Area


Image Data



Western blot analysis of lysates from NIH/3T3 and RAW264.7 cells, using ATXN2L Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from 293 cells using ATXN2L antibody.



Western Blot analysis of 293T cells using Ataxin-2L Polyclonal Antibody diluted at 1 : 2000