# **Product Name: BMP-2 Rabbit Polyclonal Antibody**

Catalog #: APRab07591



### **Summary**

BMP-2 Rabbit Polyclonal Antibody **Production Name** 

Description Rabbit Polyclonal Antibody

Host Rabbit

**Application** IF-P,IF-F,ICC/IF,WB,IHC-P,ELISA

Reactivity Human, Mouse, Rat

#### **Performance**

Conjugation Unconjugated Modification Unmodified

Isotype lgG

Clonality Polyclonal Form Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type **Buffer** 

preservative N.

**Purification** Affinity purification

#### **Immunogen**

**Gene Name** BMP2

BMP2; BMP2A; Bone morphogenetic protein 2; BMP-2; Bone morphogenetic protein **Alternative Names** 

2A; BMP-2A

Gene ID 650.0

P12643. The antiserum was produced against synthesized peptide derived from human SwissProt ID

BMP-2. AA range:226-275

### **Application**

**Dilution Ratio** 

IF-P/IF-F/ICC/IF 1:50-200, WB 1:500-1:2000, IHC-P 1:100-1:300, ELISA 1:10000.Not yet

tested in other applications.

# **Product Name: BMP-2 Rabbit Polyclonal Antibody**

Catalog #: APRab07591



Molecular Weight

60kDa


#### **Background**

This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer, which plays a role in bone and cartilage development. Duplication of a regulatory region downstream of this gene causes a form of brachydactyly characterized by a malformed index finger and second toe in human patients. [provided by RefSeq, Jul 2016],function:Induces cartilage and bone formation.,online information:Bone morphogenetic protein 2 entry,similarity:Belongs to the TGF-beta family.,subunit:Homodimer; disulfide-linked. Interacts with GREM2 (By similarity) and SOSTDC1.,tissue specificity:Particularly abundant in lung, spleen and colon and in low but significant levels in heart, brain, placenta, liver, skeletal muscle, kidney, pancreas, prostate, ovary and small intestine.,


#### Research Area

Cytokine-cytokine receptor interaction; Hedgehog; TGF-beta; Pathways in cancer; Basal cell carcinoma;

## **Image Data**




Immunohistochemistry analysis of BMP-2 antibody in paraffin-embedded human heart tissue.




Western blot analysis of lysate from HUVECcells, using BMP-2 antibody.

## **Product Name: BMP-2 Rabbit Polyclonal Antibody** Catalog #: APRab07591






Western Blot analysis of various cells using BMP-2 Polyclonal Antibody diluted at 1: 500



Western Blot analysis of HuvEc cells using BMP-2 Polyclonal Antibody diluted at 1: 500



Immunohistochemistry of paraffin-embedded human heart tissue using antibody at dilution 1:100

#### Note

For research use only.