

**Product Name: Carbonyl Reductase 1 Rabbit Polyclonal Antibody****Catalog #: APRab07921**

For research use only.

**Summary**

<b>Description</b>	Rabbit polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,IHC,ICC/IF,ELISA
<b>Reactivity</b>	Human,Rat,Mouse
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

**Application**

<b>Dilution Ratio</b>	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
<b>Molecular Weight</b>	32kDa

**Antigen Information**

<b>Gene Name</b>	CBR1 CBR1; CBR; CRN; Carbonyl reductase [NADPH] 1; 15-hydroxyprostaglandin dehydrogenase
<b>Alternative Names</b>	[NADP(+)]; NADPH-dependent carbonyl reductase 1; Prostaglandin 9-ketoreductase; Prostaglandin-E(2) 9-reductase
<b>Gene ID</b>	873.0
<b>SwissProt ID</b>	P16152
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human CBR1. AA range:181-230


## Background

The protein encoded by this gene belongs to the short-chain dehydrogenases/reductases (SDR) family, which function as NADPH-dependent oxidoreductases having wide specificity for carbonyl compounds, such as quinones, prostaglandins, and various xenobiotics. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Nov 2013], catalytic activity: (13E)-(15S)-11- $\alpha$ ,15-dihydroxy-9-oxoprost-13-enoate + NADP(+) = (13E)-11- $\alpha$ -hydroxy-9,15-dioxoprost-13-enoate + NADPH, catalytic activity: (5Z,13E)-(15S)-9- $\alpha$ ,11- $\alpha$ ,15-trihydroxyprosta-5,13-dienoate + NADP(+) = (5Z,13E)-(15S)-11- $\alpha$ ,15-dihydroxy-9-oxoprost-5,13-dienoate + NADPH, catalytic activity: R-CHOH-R' + NADP(+) = R-CO-R' + NADPH, function: Catalyzes the reduction of a wide variety of carbonyl compounds including the antitumor anthracycline antibiotics. Can convert prostaglandin E2 to prostaglandin F2- $\alpha$ , similarity: Belongs to the short-chain dehydrogenases/reductases (SDR) family, subunit: Monomer,


## Research Area

Arachidonic acid metabolism;


## Image Data



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using CBR1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, using CBR1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of 293T HELA using Carbonyl Reductase 1 Polyclonal Antibody. Antibody was diluted at 1:1000