
Product Name: CD109 Rabbit Polyclonal Antibody**Catalog #: APRab08190**

For research use only.

Summary

| | |
|----------------------|---|
| Description | Rabbit polyclonal Antibody |
| Host | Rabbit |
| Application | IHC,ICC/IF,ELISA |
| Reactivity | Human,Rat,Mouse |
| Conjugation | Unconjugated |
| Modification | Unmodified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Form | Liquid |
| Concentration | 1mg/ml |
| Storage | Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles. |
| Shipping | Ice bags |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| Purification | Affinity purification |

Application

Dilution Ratio IHC 1:50-1:200,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000

Molecular Weight

Antigen Information


| | |
|--------------------------|---|
| Gene Name | CD109 CPAMD7 |
| Alternative Names | CD109 antigen (150 kDa TGF-beta-1-binding protein;C3 and PZP-like alpha-2-macroglobulin domain-containing protein 7;Platelet-specific Gov antigen;p180;r150;CD antigen CD109) |
| Gene ID | 135228.0 |
| SwissProt ID | Q6YHK3 |
| Immunogen | Synthetic peptide from human protein at AA range: 751-800 |

Background

This gene encodes a glycosyl phosphatidylinositol (GPI)-linked glycoprotein that localizes to the surface of platelets, activated T-cells, and endothelial cells. The protein binds to and negatively regulates signalling by transforming growth factor beta (TGF-beta). Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2014],function:Modulates negatively TGFB1 signaling in keratinocytes.,polymorphism:The Gov(b) variant in position 703 defines the Gov alloantigenic determinants.,PTM:2 forms of 150 (p150) and 120 kDa (p120) exist due to proteolytic degradation from a 180 kDa form.,PTM:N-glycosylated.,similarity:Belongs to the protease inhibitor I39 (alpha-2-macroglobulin) family.,subunit:Heterodimer; disulfide-linked. Interacts with TGFB1 and TGFBR1. Forms a heteromeric complex with TGFBR1, TGFBR2 and TGFBR3 in a ligand-independent manner.,tissue specificity:Widely expressed with high level in uterus, aorta, heart, lung, trachea, placenta and in fetal heart, kidney, liver, spleen and lung. Expressed by CD34(+) acute myeloid leukemia cell lines, T-cell lines, activated T lymphoblasts, endothelial cells and activated platelets. Isoform 5 is expressed in placenta. Isoform 1 is expressed in keratinocytes and placenta.,

Research Area

Image Data



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:200