
Product Name: CD110 Rabbit Polyclonal Antibody**Catalog #: APRab08191**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight	69,40kDa

Antigen Information

Gene Name	MPL
Alternative Names	MPL; TPOR; Thrombopoietin receptor; TPO-R; Myeloproliferative leukemia protein; Proto-oncogene c-Mpl; CD110
Gene ID	4352.0
SwissProt ID	P40238
Immunogen	Synthesized peptide derived from Thrombopoietin receptor at AA range: 321-370

Background


In 1990 an oncogene, v-mpl, was identified from the murine myeloproliferative leukemia virus that was capable of

immortalizing bone marrow hematopoietic cells from different lineages. In 1992 the human homologue, named, c-mpl, was cloned. Sequence data revealed that c-mpl encoded a protein that was homologous with members of the hematopoietic receptor superfamily. Presence of anti-sense oligodeoxynucleotides of c-mpl inhibited megakaryocyte colony formation. The ligand for c-mpl, thrombopoietin, was cloned in 1994. Thrombopoietin was shown to be the major regulator of megakaryocytopoiesis and platelet formation. The protein encoded by the c-mpl gene, CD110, is a 635 amino acid transmembrane domain, with two extracellular cytokine receptor domains and two intracellular cytokine receptor box motifs . TPO-R deficient mice were severely thrombocytopenic, emphasizing the importantcaution:It is uncertain whether Met-1 or Met-8 is the initiator.,disease:Defects in MPL are a cause of congenital amegakaryocytic thrombocytopenia (CAMT) [MIM:604498]. CAMT is a disease characterized by isolated thrombocytopenia and megakaryocytopenia with no physical anomalies.,domain:The box 1 motif is required for JAK interaction and/or activation.,domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.,function:Receptor for thrombopoietin. May represent a regulatory molecule specific for TPO-R-dependent immune responses.,similarity:Belongs to the type I cytokine receptor family. Type 1 subfamily.,similarity:Contains 2 fibronectin type-III domains.,subunit:Interacts with ATXN2L.,tissue specificity:Expressed at a low level in a large number of cells of hematopoietic origin. Isoform 1 and isoform 2 are always found to be coexpressed.,


Research Area

Cytokine-cytokine receptor interaction;Jak_STAT;


Image Data



Western Blot analysis of MCF7 cells using CD110 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-liver, antibody was diluted at 1:100