Product Name: CD236 Rabbit Polyclonal Antibody
Catalog #: APRab08296
For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	IHC,ICC/IF,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	IHC 1:50-1:200,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight	

Antigen Information

Gene Name	GYPC GLPC GPC
Alternative Names	Glycophorin-C (Glycoconnectin;Glycophorin-D;GPD;Glycoprotein beta;PAS-2';Sialoglycoprotein D;CD antigen CD236)
Gene ID	2995.0
SwissProt ID	P04921
Immunogen	Synthetic peptide from human protein at AA range: 11-60

Background

Glycophorin C (GYPC) is an integral membrane glycoprotein. It is a minor species carried by human erythrocytes, but plays an

important role in regulating the mechanical stability of red cells. A number of glycophorin C mutations have been described. The Gerbich and Yus phenotypes are due to deletion of exon 3 and 2, respectively. The Webb and Duch antigens, also known as glycophorin D, result from single point mutations of the glycophorin C gene. The glycophorin C protein has very little homology with glycophorins A and B. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Feb 2012],function:This protein is a minor sialoglycoprotein in human erythrocyte membranes. The blood group Gerbich antigens and receptors for Plasmodium falciparum merozoites are most likely located within the extracellular domain. Glycophorin C plays an important role in regulating the stability of red cells.,online information:Blood group antigen gene mutation database,online information:Glycophorin C entry,polymorphism:GYPC is responsible for the Gerbich blood group system.,subcellular location:Linked to the membrane via Band 4.1.,tissue specificity:Glycophorin C is expressed in erythrocytes. Glycophorin D is ubiquitous.,

Research Area

Cardiovascular

Image Data



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-spleen, antibody was diluted at 1:200