
Product Name: CEP57 Rabbit Polyclonal Antibody**Catalog #: APRab08668**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:20000-1:40000
Molecular Weight	50kDa

Antigen Information


Gene Name	CEP57
Alternative Names	CEP57; KIAA0092; TSP57; Centrosomal protein of 57 kDa; Cep57; FGF2-interacting protein; Testis-specific protein 57; Translokin
Gene ID	9702.0
SwissProt ID	Q86XR8
Immunogen	The antiserum was produced against synthesized peptide derived from human CEP57. AA range:241-290

Background

This gene encodes a cytoplasmic protein called Translokin. This protein localizes to the centrosome and has a function in microtubular stabilization. The N-terminal half of this protein is required for its centrosome localization and for its multimerization, and the C-terminal half is required for nucleating, bundling and anchoring microtubules to the centrosomes. This protein specifically interacts with fibroblast growth factor 2 (FGF2), sorting nexin 6, Ran-binding protein M and the kinesins KIF3A and KIF3B, and thus mediates the nuclear translocation and mitogenic activity of the FGF2. It also interacts with cyclin D1 and controls nucleocytoplasmic distribution of the cyclin D1 in quiescent cells. This protein is crucial for maintaining correct chromosomal number during cell division. Mutations in this gene cause mosaic variegated aneuploidy syndrome, a rare autosomal recessive disorder. Multiplefunction:Mediates nuclear translocation and mitogenic activity of the internalized growth factor FGF2.,similarity:Belongs to the translokin family.,subcellular location:Associates with microtubules and the centrosome.,subunit:Homodimer. Interacts with FGF2 and RAP80. Does not interact with FGF1 or FGF2 isoform 24 kDa.,tissue specificity:Ubiquitous.,

Research Area

Image Data



Western blot analysis of lysates from COLO and LOVO cells, using CEP57 Antibody. The lane on the right is blocked with the synthesized peptide.