
Product Name: CNOT4 Rabbit Polyclonal Antibody**Catalog #: APRab09129**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
Molecular Weight	64kDa

Antigen Information

Gene Name	CNOT4
Alternative Names	CNOT4; NOT4; CCR4-NOT transcription complex subunit 4; CCR4-associated factor 4; E3 ubiquitin-protein ligase CNOT4; Potential transcriptional repressor NOT4Hp
Gene ID	4850.0
SwissProt ID	O95628
Immunogen	The antiserum was produced against synthesized peptide derived from human CNOT4. AA range:61-110


Background

CCR4-NOT transcription complex subunit 4(CNOT4) Homo sapiens The protein encoded by this gene is a subunit of the CCR4-NOT complex, a global transcriptional regulator. The encoded protein interacts with CNOT1 and has E3 ubiquitin ligase activity. Several transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jul 2010],function:Has E3 ubiquitin ligase activity. The CCR4-NOT complex functions as general transcription regulation complex.,pathway:Protein modification; protein ubiquitination.,PTM:Autoubiquitinated.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,sequence caution:Sequencing errors.,similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 RING-type zinc finger.,similarity:Contains 1 RRM (RNA recognition motif) domain.,subunit:Subunit of the CCR4-NOT core complex that contains CHAF1A, CHAF1B, CNOT1, CNOT2, CNOT3, CNOT4, CNOT6 and CNOT8. Binds CNOT1 via its C-terminus. Binds E2 ubiquitin ligases via its RING domain.,


Research Area

RNA degradation;

Image Data



Immunohistochemistry analysis of paraffin-embedded human testis tissue, using CNOT4 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat and K562 cells, using CNOT4 Antibody. The lane on the right is blocked with the synthesized peptide.