## **Product Name: COL14A1 Rabbit Polyclonal Antibody**

Catalog #: APRab09170



#### **Summary**

Production Name COL14A1 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

**Host** Rabbit

**Application** IHC-P,IF-P,IF-F,ICC/IF,ELISA

**Reactivity** Human, Mouse, Rat

#### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

Clonality Polyclonal Form Liquid

**Storage** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type

preservative N.

**Purification** Affinity purification

#### **Immunogen**

Buffer

Gene Name COL14A1 UND

Alternative Names Collagen alpha-1(XIV) chain (Undulin)

**Gene ID** 7373.0

**SwissProt ID** Q05707. Synthetic peptide from human c-ternal protein at AA range: 1571-1620

### **Application**

**Dilution Ratio** IHC-P 1:50-200, ELISA 1:10000-20000, IF-P/IF-F/ICC/IF 1:50-200

**Molecular Weight** 

#### **Background**

This gene encodes the alpha chain of type XIV collagen, a member of the FACIT (fibril-associated collagens with interrupted


# Product Name: COL14A1 Rabbit Polyclonal Antibody Catalog #: APRab09170



triple helices) collagen family. Type XIV collagen interacts with the fibril surface and is involved in the regulation of fibrillogenesis. [provided by RefSeq, Jan 2013], function: Plays an adhesive role by integrating collagen bundles. It is probably associated with the surface of interstitial collagen fibrils via COL1. The COL2 domain may then serve as a rigid arm which sticks out from the fibril and protrudes the large N-terminal globular domain into the extracellular space, where it might interact with other matrix molecules or cell surface receptors., PTM: Lysines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in all cases and bind carbohydrates., PTM: May contain numerous cysteine residues involved in inter- and intramolecular disulfide bonding., PTM: Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains., similarity: Belongs to the fibril-associated collagens with interrupted helices (FACIT) family., similarity: Contains 1 TSP N-terminal (TSPN) domain., similarity: Contains 2 VWFA domains., similarity: Contains 8 fibronectin type-III domains., subunit: Homotrimer.,

#### **Research Area**

#### **Image Data**



Immunohistochemical analysis of paraffin-embedded human-skin, antibody was diluted at 1:200

#### Note

For research use only.