
Product Name: COX11 Rabbit Polyclonal Antibody**Catalog #: APRab09266**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000
Molecular Weight	31kDa

Antigen Information

Gene Name	COX11
Alternative Names	COX11; Cytochrome c oxidase assembly protein COX11; mitochondrial
Gene ID	1353.0
SwissProt ID	Q9Y6N1
Immunogen	The antiserum was produced against synthesized peptide derived from human COX11. AA range:51-100

Background


Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer

from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes a protein which is not a structural subunit, but may be a heme A biosynthetic enzyme involved in COX formation, according to the yeast mutant studies. However, the studies in *Rhodobacter sphaeroides* suggest that this gene is not required for heme A biosynthesis, but required for stable formation of the Cu(B) and magnesium centers of COX. This human protein is predicted to function: Exerts its effect at some terminal stage of cytochrome c oxidase synthesis, probably by being involved in the insertion of the copper B into subunit I., similarity: Belongs to the COX11/ctaG family., subunit: Interacts with CNNM4/ACDP4., tissue specificity: Ubiquitous.,

Research Area

Oxidative phosphorylation;

Image Data



Western blot analysis of lysates from RAW264.7 cells, using COX11 Antibody. The lane on the right is blocked with the synthesized peptide.