
Product Name: CP4FN Rabbit Polyclonal Antibody**Catalog #: APRab09304**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Rat,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,ELISA 1:5000-1:20000
Molecular Weight	58kDa

Antigen Information

Gene Name	CYP4F22
Alternative Names	
Gene ID	126410.0
SwissProt ID	Q6NT55
Immunogen	Synthesized peptide derived from human protein . at AA range: 440-520


Background

cytochrome P450 family 4 subfamily F member 22(CYP4F22) Homo sapiens This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This gene is part of a cluster of cytochrome P450 genes

on chromosome 19 and encodes an enzyme thought to play a role in the 12(R)-lipoxygenase pathway. Mutations in this gene are the cause of ichthyosis lamellar type 3. [provided by RefSeq, Jul 2008],cofactor:Heme group.,disease:Defects in CYP4F22 are the cause of ichthyosis lamellar type 3 (LI3) [MIM:604777]. LI is a non-bullous ichthyosis, a skin disorder characterized by abnormal cornification of the epidermis. It is one the most severe forms of ichthyoses apparent at birth and persisting throughout life. LI patients are born encased in a tight, shiny, translucent covering called collodion membrane. Over the first weeks of life, the collodion membrane is gradually replaced by generalized large, dark brown, plate-like scales with minimal to no erythroderma. Tautness of facial skin commonly results in ectropion, eclabium and scarring alopecia of the scalp. Common complications are severe heat intolerance and recurrent ear infections.,similarity:Belongs to the cytochrome P450 family.,

Research Area

Image Data



Western Blot analysis of HEK293 lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000