
Product Name: Creatine Kinase M Rabbit Polyclonal Antibody**Catalog #: APRab09373**

For research use only.

Summary

Description	Rabbit polyclonal Antibody
Host	Rabbit
Application	WB,IHC,ICC/IF,ELISA
Reactivity	Human,Mouse,Rat
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Concentration	1mg/ml
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
Purification	Affinity purification

Application

Dilution Ratio	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:20000-1:40000
Molecular Weight	43kDa

Antigen Information

Gene Name	CKM
Alternative Names	CKM; CKMM; Creatine kinase M-type; Creatine kinase M chain; M-CK
Gene ID	1158.0
SwissProt ID	P06732
Immunogen	The antiserum was produced against synthesized peptide derived from human M-CK. AA range:10-59

Background


The protein encoded by this gene is a cytoplasmic enzyme involved in energy homeostasis and is an important serum marker

for myocardial infarction. The encoded protein reversibly catalyzes the transfer of phosphate between ATP and various phosphogens such as creatine phosphate. It acts as a homodimer in striated muscle as well as in other tissues, and as a heterodimer with a similar brain isozyme in heart. The encoded protein is a member of the ATP:guanido phosphotransferase protein family. [provided by RefSeq, Jul 2008],catalytic activity:ATP + creatine = ADP + phosphocreatine.,function:Reversibly catalyzes the transfer of phosphate between ATP and various phosphogens (e.g. creatine phosphate). Creatine kinase isoenzymes play a central role in energy transduction in tissues with large, fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa.,online information:CKM entry,online information:Creatine kinase entry,similarity:Belongs to the ATP:guanido phosphotransferase family.,subunit:Dimer of identical or non-identical chains. With MM being the major form in skeletal muscle and myocardium, MB existing in myocardium, and BB existing in many tissues, especially brain.,


Research Area

Arginine and proline metabolism;


Image Data




Immunofluorescence analysis of HepG2 cells, using M-CK Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human heart tissue, using M-CK Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from Jurkat, HeLa, HepG2, and HUVEC cells, using M-CK Antibody. The lane on the right is blocked with the synthesized peptide.




Western blot analysis of the lysates from HepG2 cells using M-CK antibody.



Western Blot analysis of various cells using Creatine Kinase M Polyclonal Antibody



Western Blot analysis of Jurkat cells using Creatine Kinase M Polyclonal Antibody



The picture was kindly provided by our customer