

**Product Name: CSRNP3 Rabbit Polyclonal Antibody****Catalog #: APRab09468**

For research use only.

**Summary**

<b>Description</b>	Rabbit polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human,Mouse,Monkey
<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Concentration</b>	1mg/ml
<b>Storage</b>	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
<b>Shipping</b>	Ice bags
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

**Application**

<b>Dilution Ratio</b>	WB 1:500-1:2000,ELISA 1:10000-1:20000
<b>Molecular Weight</b>	64kDa

**Antigen Information**


<b>Gene Name</b>	CSRNP3
<b>Alternative Names</b>	CSRNP3; FAM130A2; TAIP2; Cysteine/serine-rich nuclear protein 3; CSRNP-3; Protein FAM130A2; TGF-beta-induced apoptosis protein 2; TAIP-2
<b>Gene ID</b>	80034.0
<b>SwissProt ID</b>	Q8WYN3
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human TAIP-2. AA range:171-220

**Background**

function: Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional activator activity. Plays a role in apoptosis., sequence caution: Unusual initiator. The initiator methionine is coded by a non-canonical CTG leucine codon., similarity: Belongs to the AXUD1 family., function: Binds to the consensus sequence 5'-AGAGTG-3' and has transcriptional activator activity. Plays a role in apoptosis., sequence caution: Unusual initiator. The initiator methionine is coded by a non-canonical CTG leucine codon., similarity: Belongs to the AXUD1 family.,

## Research Area

## Image Data



Western blot analysis of the lysates from HepG2 cells using TAIP-2 antibody.